A Suns Water Study and Pre-Publication - Resolving the Coronal Heating Paradox

This is a comprehensive and detailed summary of the scientific article and short study about one of the biggest unsolved physical problems. This preview and publication here was not completly corrected and formatted, this will be done later after some further updates. It is an extract and preview for the final publication. It is also an introduction for a new book with focus on scientific issues related to solar science.

Solving Problems of Solar Physics: Evidence, Key Factors and Mechanisms

The coronal heating paradox refers to the fundamental mystery of why the Sun's corona, the outermost layer of the solar atmosphere, is vastly hotter than the photosphere beneath it. Despite being farther from the Sun's energy-producing core, the corona reaches temperatures of 1-3 million Kelvin, whereas the photosphere is a relatively cool and can heat up to 5,500°C (5,800 K). The scientific paradox challenges traditional models of solar energy transport and has led to intense research into the underlying mechanisms that explain the temperature difference. Recent developments in solar physics, including studies of magnetic field dynamics, wave-particle interactions, and particle acceleration, have provided comprehensive solutions to this puzzle.

During studies for the Sun's Water Theory the lead researcher discovered many solution approaches to known problems of solar science. He found many crucial factors and answers to resolve the coronal paradox almost completely – together with known and unknown solutions. It was one of the most perplexing problems in solar physics, first identified in the 1940s, where observations show that the temperature of the Sun's outer atmosphere is much higher than its visible surface or photosphere with an average temperature of around 5,500 K. The temperature discrepancy has been a longstanding mystery with the central question: How can the Sun's corona be so much hotter than the photosphere? The main question is answered and many of the problems are solved in this scientific article and in other sections of the special Suns Water study project with further research. It follows a comprehenisve overview of main factors that can drive coronal heating.

1. Alfvén Waves and Their Role in Heating Processes

Alfvén waves are primary candidates for transporting energy from the Sun's interior to the corona. These magnetohydrodynamic (MHD) waves propagate along magnetic field lines in plasma. The dispersion relation for Alfvén waves is: $\omega^2 = k^2 v_A^2$, where ω is the wave frequency, k is the wave number, and v_A is the Alfvén speed ($v_A = B/\sqrt{(\mu_0 \rho)}$), which depends on the magnetic field strength B, plasma density ρ, and permeability of free space μ_0. These waves are widely discussed as contributors to coronal heating. They transport both energy and momentum, and when dissipated, transfer energy to the plasma, causing heating. A key feature of Alfvén waves in coronal heating is their ability to carry energy from the solar surface (photosphere) to the corona. The dissipation of Alfvén waves involves resonant absorption, where wave energy is absorbed by plasma at specific frequencies. This process depends on magnetic field geometry, plasma density, and wave frequency.

A modified model for resonant absorption can be written as: $dE/dt = (B^2 / \mu_0) * [v_r / (1 + v_r^2)]$, where **dE/dt** is the rate of energy absorption, **B** is the magnetic field strength, **μ_0** is the permeability of free space, and **v_r** is the wave frequency relative to the plasma's resonant frequency. For temperature variations, the speed is modified as: **$v_A = B / \sqrt{(\mu_0 \rho)} * (1 + \delta T)$** , where **$\delta$** is a constant and **$T$** is the temperature (K). This modification accounts for the temperature dependence of wave propagation. In high-temperature plasmas, resistivity changes influence wave dissipation rates. Temperature fluctuations, such as during solar flares and **coronal mass ejections (CMEs)**, can enhance wave heating. Observations from Hinode and SDO have detected wave modes in coronal loops, supporting resonant absorption with frequencies that match theoretical predictions. Studies of the Sun's Water Theory will refine these calculations, aiding the understanding of solar processes.

A further modification for wave energy dissipation is: $v_A = (B + E_w) / \sqrt{(\mu_0 \rho)}$, where **E_w** is the energy contribution from Alfvén waves. This adjustment reflects the significant role of wave dissipation in heating the corona. Alfvén wave energy interacts with plasma, potentially causing nonlinear wave processes that deposit energy into the plasma. Wave damping occurs through mechanisms like resonant absorption and mode conversion. Understanding how energy is transferred and dissipated is crucial. Turbulent cascading in the corona can enhance wave dissipation. As small-scale turbulence develops, Alfvén

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

waves transfer energy across scales, converting it into heat. This process may explain the high temperatures in the corona relative to the photosphere, providing continuous energy input to the coronal plasma. More details on Alfvén waves and heating mechanisms can be found in other sections.

Observational and Empirical Evidence: The role of Alfvén waves in coronal heating has been examined using high-resolution instruments like **Hinode, SDO, and TRACE**. Observations of coronal loops and active regions have revealed oscillations matching the signatures of Alfvén waves. Specifically, **SDO's Atmospheric Imaging Assembly (AIA)** detected high-frequency oscillations in coronal loops, with periods corresponding to Alfvén wave frequencies. These waves are believed to propagate upward from the photosphere, carrying energy to the corona. **Hinode's EUV Imaging Spectrometer (EIS)** has observed resonant absorption in coronal loops. Studies of quasi-periodic pulsations (QPPs) in flare regions also support the presence of Alfvén waves and their role in energy dissipation. Furthermore, coronal seismology with SDO data has confirmed that observed oscillations align with theoretical Alfvén wave predictions. These waves contribute to localized heating and are thought to play a major role in the high temperatures observed in the corona.

2. Bow Shocks and Bernoulli's Equation

Bernoulli's equation and bow shocks provide insights into the coronal heating paradox, especially regarding energy transfer, plasma dynamics, and solar wind acceleration. While they don't fully resolve the paradox, these concepts shed light on specific processes related to the heating and acceleration of the solar corona. Bow Shocks and Energy Dissipation Bow shocks occur when the solar wind interacts with surrounding media, such as the Earth's magnetosphere or interstellar space. While typically associated with planetary magnetospheres, the concept also helps understand shock heating in the Sun's corona, particularly in active regions and where the solar wind interacts with the Sun's magnetic fields. In the corona, shock waves, generated by reconnection or plasma flows, can dissipate energy as magnetic fields reorganize. These shocks convert the kinetic energy of plasma into thermal energy, contributing to coronal heating.

Shock Heating and Plasma Dynamics: When the solar wind encounters denser or slower-moving regions (e.g., near active regions or coronal loops), bow shocks compress and heat the plasma. The interaction between fast solar wind and slower coronal material creates a shock front, dissipating energy as heat, which increases the temperature of the corona. **Shock Heating Equation: $\Delta T = (\gamma - 1) * (P2 - P1) / \rho$** , where **$\Delta T$** is the temperature increase across the shock, **γ** is the adiabatic index (specific heat ratio), **P2** and **P1** are the pressures before and after the shock, and **ρ** is the plasma density. This equation shows how the shock dissipates the kinetic energy of solar wind particles, converting it into heat, potentially explaining the temperature differences observed in the corona.

Bernoulli's equation relates the pressure, velocity, and potential energy of a fluid in steady flow. Although it is primarily used for incompressible fluids, its principles can be adapted to the solar corona to understand the energy balance between coronal heating and solar wind acceleration. The equation is expressed as: **$P + 0.5 * \rho * v^2 + \rho * g * h = \text{constant}$** , where **P** is pressure, **ρ** is plasma density, **v** is plasma velocity, **g** is gravitational acceleration, and **h** is height. In the corona, this equation helps link the pressure and velocity of the solar wind, illustrating the connection between coronal heating and solar wind acceleration. As plasma in the corona heats, it expands and accelerates, moving from high-pressure, high-temperature regions to lower-pressure areas in the outer corona and solar wind.

Bernoulli's Energy Balance in the Solar Wind: $\Delta P = 0.5 * \rho * (v2^2 - v1^2)$, where **ΔP** is the pressure difference between two points, **ρ** is plasma density, and **v2** and **v1** are the velocities at two different points in the corona or solar wind. This equation shows that an increase in velocity leads to a pressure decrease, which aligns with observations that solar wind accelerates as it moves away from the Sun's corona. The relationship between thermal, kinetic, and magnetic energy in the corona may influence coronal heating, providing insight into energy transfer mechanisms.

Pressure-Velocity Relationship in the Corona: The dissipation of magnetic energy (e.g., through magnetic reconnection) could increase the corona's pressure. According to Bernoulli's principle, this could cause a corresponding increase in plasma velocity. The accelerated plasma then carries energy away from the corona, contributing to both coronal heating and the outward flow of solar wind. In regions with higher solar wind speed, Bernoulli's equation suggests that thermal energy is converted into kinetic energy, driving

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

solar wind acceleration. Therefore, magnetic energy conversion to heat in the corona may also drive solar wind acceleration, linking heating and wind processes.

Bow shocks and Bernoulli's equation together offer explanations for solar wind acceleration. Both mechanisms transfer energy from thermal or magnetic forms into kinetic energy as plasma moves outward. Bow shocks dissipate energy from turbulent or fast solar wind, increasing plasma temperature in localized regions. In areas where the solar wind speed is low and coronal pressure is high, Bernoulli's equation suggests thermal energy conversion into kinetic energy, driving solar wind acceleration. Therefore, energy conversion at shock fronts and velocity transitions may be responsible for both coronal heating and solar wind acceleration.

Coronal Heating through Shock-Driven Waves: Shock-driven waves, generated by turbulence or reconnection events, could propagate through the corona, similar to how bow shocks dissipate energy in planetary magnetospheres. These waves could contribute to coronal heating by converting kinetic energy into thermal energy. The propagation of waves (e.g., Alfvén waves) through the corona leads to energy dissipation when they encounter plasma regions with varying densities or magnetic fields, creating localized heating. These shock- and wave-driven mechanisms might help explain how small-scale processes like nanoflares or magnetic reconnection sustain the high temperature in the solar atmosphere, even though the source of this heat remains unclear.
While bow shocks and Bernoulli's equation do not provide definitive solutions to the coronal heating paradox, they offer valuable insights, especially regarding energy transfer and plasma dynamics in the corona and solar wind. These concepts help explain how thermal energy is converted to kinetic energy, contributing to solar wind acceleration, and how magnetic energy dissipation might lead to localized coronal heating. Further research into how these mechanisms interact with other processes, such as wave dissipation and turbulence, will deepen our understanding of coronal heating.

3. Chemical Processes and Element Abundance

The Sun's corona, primarily composed of plasma, plays a crucial role in energy transport and magnetic structure formation, which influence heating processes. While the coronal heating paradox is often framed in terms of physical mechanisms such as waves, reconnection, and turbulence, chemical processes and element abundances are also vital factors in determining the observed temperature distribution in the corona. The chemical composition, dominated by hydrogen and helium, includes trace amounts of heavier elements such as oxygen, carbon, neon, iron, and silicon. The interactions between these elements significantly influence both heating and cooling rates in the corona.

Hydrogen is the most abundant element in the solar wind and corona. It primarily exists as protons (H^+), which are crucial in maintaining the ionization equilibrium, especially at temperatures exceeding 1 million K. Proton interactions with the surrounding plasma through Coulomb collisions with electrons and heavier ions facilitate energy transfer, thereby heating the electron population, which in turn heats the ions. Proton-ion and proton-electron collisions contribute to energy dissipation and radiative losses via **Bremsstrahlung**. The low ionization potential of hydrogen (~13.6 eV) makes it easily ionizable in the corona, and as a result, **H^+ ions** participate significantly in energy absorption during magnetic reconnection events and wave-particle interactions.

Contributions of Specific Elements to Coronal Heating: Carbon (C), especially in its **C^{4+} state**, plays an important role in coronal heating. C^{4+} ions are highly effective at absorbing UV radiation, thus maintaining the ionization balance in the corona. Carbon ions contribute to radiative cooling through emission lines in the UV and X-ray ranges, particularly the **carbon Ly-α** line. As the corona's temperature increases, carbon ions transition through various ionization states, which significantly impacts the thermal equilibrium and energy transport in the corona. Their interaction with the plasma through electron-ion collisions also contributes to energy dissipation.

Helium (He), primarily in the forms of **He^+** and **He^{2+}**, contributes to the radiative cooling of the corona. Although less abundant than hydrogen, helium's role remains significant due to its emission and absorption processes. The **He^{2+}** ion, with a higher ionization energy, interacts with high-energy photons and charged particles, particularly during magnetic field interactions and wave heating. These interactions lead to the

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

emission of UV radiation, which plays a role in the overall cooling of the corona. Helium's interactions with the magnetic field are also crucial in regulating the ionization balance of the plasma.

Iron (Fe), one of the most chemically significant elements in the solar corona, exists in highly ionized states (**Fe^{6+}, Fe^{7+}, Fe^{8+}, etc.**). Iron ions are vital for both the absorption and emission of radiation. **Fe^{6+}** ions, in particular, absorb high-energy UV and X-ray radiation and re-emit energy in the form of **X-ray lines**, contributing to significant radiative losses. These ions are found in the hottest regions of the corona, where they also contribute to **Bremsstrahlung radiation**, further enhancing energy dissipation. The high charge states of iron ions allow them to interact significantly with both electromagnetic radiation and magnetic fields, especially during solar flares or **coronal mass ejections (CMEs)**.

Iron's Role in Magnetic Reconnection and Wave Heating: Iron ions, especially **Fe^{6+}** and **Fe^{7+}**, interact with magnetic fields and Alfvén waves in the corona. These ions efficiently absorb energy from MHD waves and magnetic reconnection processes, which leads to ion acceleration, increased ion temperature, and subsequent heating of the plasma. During **solar flares** or **CMEs**, these interactions can contribute to magnetic energy dissipation and ion heating, thereby enhancing the coronal heating process.

Magnesium (Mg), primarily in the form of **Mg^{9+}** and **Mg^{7+}** ions, contributes to energy dissipation through **collisional ionization** and **Bremsstrahlung processes**. Although less abundant than iron or oxygen, magnesium plays an important role in wave-particle interactions, especially at higher temperatures. Neon (Ne), in the form of **Ne^{6+}** and **Ne^{7+}**, also plays a role in coronal heating by absorbing UV radiation and emitting X-rays. Neon's ability to maintain a high ionization state at the temperatures present in the corona makes it an efficient contributor to the thermal balance of the plasma.

Nitrogen (N): Although less abundant, nitrogen, especially in its **N^{4+}** state, plays a role in radiative cooling. It absorbs high-energy radiation and emits UV and X-rays when excited by energetic events such as solar flares or magnetic reconnection. While its contribution to the overall coronal heating process is smaller than other elements like carbon or oxygen, nitrogen still impacts the energy dynamics, particularly in the **transition region** between the photosphere and the corona.

Oxygen (O) is one of the most important elements in the solar corona, primarily in the form of **O^{6+}** and **O^{7+}** ions. Oxygen ions are abundant in the transition region and upper corona, where they absorb and emit radiation in the UV and X-ray spectra. **O^{6+}** ions, in particular, absorb significant amounts of solar radiation, and undergo recombination and excitation processes that contribute to **radiative losses**. Oxygen also plays a role in **energy dissipation** through **Bremsstrahlung** and collisional ionization, with **O^{6+}** ions being highly sensitive to changes in electron density and temperature, providing critical information about coronal heating mechanisms.

Silicon (Si), in its highly ionized states (**Si^{6+}** and **Si^{7+}**), contributes to both radiative cooling and ionization balance in the corona. Like iron, silicon emits radiation in the X-ray range and absorbs energy from magnetic fields and waves, releasing it as thermal radiation. Silicon ions contribute to the overall heating processes by interacting with energetic solar events, such as flares or CMEs, and dissipating energy into the surrounding plasma.

Collisional Excitation and De-excitation: Collisions between electrons and ions lead to the excitation of ions to higher energy states, and as the ions relax to lower energy states, they emit photons. This process, known as **collisional de-excitation**, can be described by the rate: **$R_{exc} = n_e * n_i * \sigma_{exc} * v_{rel}$** , where **$\sigma_{exc}$** is the excitation cross-section, and **v_{rel}** is the relative velocity between electrons and ions. The emitted photons, observed as spectral lines in the UV and X-ray ranges, provide essential diagnostic information about the temperature and density of the plasma. These emission lines are crucial for diagnosing energy transfer and dissipation mechanisms in the corona. The rate of **collisional ionization** can be expressed as: **$R_{ci} = n_e * n_i * \sigma_{ci} * v_{rel}$** , where **$\sigma_{ci}$** is the collisional ionization cross-section, and this process helps maintain the ionization state in the corona, especially in hotter regions.

Ionization and Recombination Processes: Ionization rates are crucial for understanding the dynamics of energy transfer in the corona. The ionization of elements like hydrogen, oxygen, and iron influences **plasma conductivity** and the overall energy balance. The **Saha equation** is often used to model ionization equilibria in stellar atmospheres: **$(n_i / n_e) = (2 * \pi * m_e * k_B * T)^{(3/2)} * (g_i / g_e) * \exp(-I_i / k_B T)$** , where **$n_i$** is the ion density, **n_e** the electron density, **T** is temperature, and **I_i** is the ionization energy.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

The balance between ionization and recombination rates dictates the ionization state of the plasma, influencing coronal heating processes. Recombination, the reverse of ionization, occurs when a free electron combines with an ion to form a neutral species. The **recombination rate** is the inverse of the ionization rate, and the rate of recombination is given by: **$R_{recomb} = n_e * n_i * \beta(T)$** , where **$\beta(T)$** is the recombination coefficient, dependent on the plasma temperature. The interplay between ionization and recombination rates is essential for maintaining the ionization balance, and thus for controlling the energy dynamics in the corona.

Line Emission and Ionized Metals: Ionized metals like **Fe**, **O**, **C**, and **Si** contribute to coronal heating through line emissions, which provide diagnostic information about the plasma's temperature and ionization state. These emissions arise from **collisional excitation** followed by **de-excitation** and are observed in solar spectroscopy. The line intensity, given by: **$L_{line} = n_e * n_i * \sigma_{exc} * h * \nu$** , where **h** is Planck's constant and **ν** is the frequency of the emitted radiation, helps determine the plasma's thermal state and is key to understanding energy dissipation in the corona.

High-Temperature Chemistry: The ionization states of elements like iron, magnesium, and silicon can affect the thermal conductivity of the plasma and influence wave energy dissipation. The ionization and recombination processes are temperature-dependent, and they modify the plasma's heat capacity. The overall state of the plasma is described by the equation of state (EOS), which links temperature, density,

4. Convection and Energy Transport in the Photosphere

Convection is a crucial mechanism of energy transport in the Sun's convective zone, just below the photosphere. In this zone, thermal convection dominates: hot plasma rises, cools as it nears the surface, and sinks again. This convective energy transport takes over when radiation alone is no longer sufficient to transfer energy through the Sun's outer layers.

Convective Energy Transport (Adiabatic Process): At around 0.7 solar radii, the Sun's temperature drops to about 2 million K, and the plasma becomes opaque to radiation. This is when convection becomes the primary energy transport mechanism. The convective zone extends from 0.7 to 1.0 solar radii and involves the rise of hot plasma and sinking of cooler plasma, akin to a boiling pot of water. In this region, convection becomes dominant because conduction is inefficient due to the opacity of the material.
The convective heat flux is given by the equation: **$Q = (dT/dz) * c_p * \rho * \nu$** , where **dT/dz** is the temperature gradient, **c_p** the specific heat capacity at constant pressure, **ρ** the plasma density and **ν** is the convective velocity. This equation shows how the temperature gradient drives the convection process and transports energy from the deeper layers of the Sun to the photosphere. The convective motion is driven by buoyancy: when plasma at lower levels heats up, it becomes less dense and rises. Upon reaching the surface, it cools, becomes denser, and sinks, continuing this cycle. The convection process forms granules and supergranules on the Sun's surface, these small convection cells (~1000 km in size) are visible in solar observations. The convection cells are caused by the decreasing temperature with altitude, which triggers hot material to rise and cooler material to sink, creating a dynamic convection pattern.

Convective Velocity and Granulation: The Rayleigh number helps describe the transition from stable conduction to turbulent convection. When Ra exceeds a certain threshold, convection becomes turbulent, resulting in the formation of large convection cells. The granules seen on the Sun's surface are part of these cells. The Rayleigh number is given by: **$Ra = (g * \beta * \Delta T * L^3) / (\nu * \alpha)$** , where **g** is the gravitational acceleration, **β** the thermal expansion coefficient, **ΔT** the temperature difference, **L** the characteristic length scale (the convective cell size), **ν** the kinematic viscosity and **α** is the thermal diffusivity. Once the energy reaches the photosphere, it is radiated as electromagnetic radiation, including visible light and ultraviolet radiation. The photosphere marks the point where the plasma becomes transparent enough for light to escape. The temperature here is approximately 5,500 K, and photons can travel freely from this layer into space.

5. Energy Conservation and Dissipation

Understanding the energy conservation and dissipation mechanisms in the corona is key to addressing the coronal heating paradox. Despite the corona being much hotter than the photosphere, traditional energy

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

transport mechanisms like radiation and conduction cannot fully explain this temperature difference. A comprehensive analysis of energy conservation and dissipation is therefore necessary.

Cooling by Radiation: A major cooling mechanism in the corona is Bremsstrahlung radiation, which occurs when charged particles, such as electrons, are decelerated or accelerated in the presence of other charged particles (ions). The cooling rate due to Bremsstrahlung is given by: $L = n_e^2 \Lambda(T)$, where n_e is the electron number density and $\Lambda(T)$ is the radiative loss function, which depends on the temperature of the plasma. This form of cooling is most significant in the lower corona, where the plasma density is higher and radiation losses are more prominent.

Dissipation of Waves in the Corona: Alfvén waves are a key mechanism for transporting energy from the photosphere to the corona. These waves must dissipate their energy to heat the plasma in the corona. This can happen via several mechanisms, including mode conversion (where Alfvén waves turn into magnetoacoustic waves) and resonant absorption (where the wave energy is absorbed by the plasma). The heating rate from wave dissipation can be expressed as: $P_{wave} = \alpha W$, where α is the wave dissipation efficiency (a dimensionless constant) and W is the energy carried by the wave. Once the wave energy is dissipated, it is transferred to the plasma, heating it.

Energy Balance Equation for the Corona: In a steady-state system, the energy balance equation for the solar corona can be written as: $(\partial E/\partial t) + \nabla * F = Q - L$, where E is the total energy density, F the energy flux, Q the heating rate (which includes mechanisms like wave dissipation and magnetic reconnection) and L is the cooling rate (due to radiation and conduction). This equation captures how energy is transported, supplied by heating mechanisms, and lost through cooling processes.

Energy Flux and Heat Conduction: The energy flux is described by Fourier's Law: $F = -\kappa \nabla T$, where: κ is the thermal conductivity and ∇T is the temperature gradient.
In the solar corona, the Spitzer conductivity is used to model heat transport: $\kappa = 1.84 \times 10^{-5} T^{(5/2)} / n_e$, where T is the temperature and n_e is the electron number density. In the presence of magnetic fields, thermal conduction is suppressed perpendicular to the field lines. The effective conductivity in the corona is much lower than the Spitzer value for an unmagnetized plasma. This reduced thermal conductivity may contribute to the build-up of heat in magnetic loops, helping to explain why the corona is hotter than the photosphere.

Heat Loss via Expansion (Adiabatic Cooling): Another important cooling mechanism in the corona is adiabatic cooling. As plasma moves outward in the solar wind, its density decreases, causing an expansion that leads to cooling. This cooling becomes more significant at higher altitudes, where plasma expands more rapidly. The cooling rate due to expansion is: $L_{expansion} = n_e T / (\gamma - 1) \nabla v$, where γ is the adiabatic index (about 5/3 for a monoatomic ideal gas) and v is the plasma velocity.

Refined Energy Balance for the Corona: While the basic energy balance equation captures the general idea of heating and cooling in the corona, a more refined model must consider complex processes like turbulent energy cascades, resistive dissipation, and wave-particle interactions. A more detailed energy balance equation that includes these dissipative processes is: $(\partial E/\partial t) + \nabla * (\kappa \nabla T) = P_{source} - P_{loss}$, where P_{source} is the heating rate from mechanisms such as wave dissipation, magnetic reconnection, and non-thermal particle acceleration. P_{loss} is the cooling rate, which accounts for radiative cooling, adiabatic cooling, turbulent dissipation, and resistive effects. This refined equation demonstrates that energy is not only transported via conduction but is also generated by heating mechanisms and lost through various cooling processes. Improving our understanding of energy input (such as wave dissipation) and refining the loss mechanisms (such as turbulent dissipation) will help explain the high temperatures observed in the corona.

Energy Losses and Cooling Mechanisms: One of the challenges in the coronal heating paradox is understanding how energy is stored and dissipated in the corona despite substantial energy losses. Radiation losses, especially in regions with high density and temperature, are dominated by optically thin emission processes from ionized ions like Fe IX-XIII. The radiative cooling rate can be expressed as: $Q_{rad} = n_e * n_i * \Lambda(T)$, where n_e is the electron density, n_i is the ion density and (T) is the radiative cooling function. In the corona, the presence of strong magnetic fields limits the efficiency of thermal conduction perpendicular to the field lines. The Spitzer conductivity for an unmagnetized plasma can be used. However, in the corona, the effective conductivity is much lower due to the magnetic fields, which hinder the flow

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

of heat perpendicular to the magnetic field lines. This lower thermal conductivity suggests that heat cannot efficiently escape along magnetic loops, allowing thermal energy to build up, thus explaining the higher temperatures in the corona compared to the photosphere.

6. Energy Transport in the Sun's Layers

Energy transport in the Sun's layers varies significantly with depth and plasma conditions. In the photosphere, radiation dominates, although convective upwelling creates granulation - regions of rising hot plasma surrounded by cooler sinking material. This convection pattern visibly transports energy at the Sun's surface. In the outer layers (chromosphere and corona) energy transport becomes more complex. The chromosphere still involves radiation but is more transparent at lower temperatures. In the corona, energy is primarily transported via magnetic reconnection, Alfvén waves, and turbulent processes. This increase in temperature with height is central to the coronal heating paradox, where mechanisms like wave dissipation and magnetic dynamics help explain the high corona temperatures. The Sun's energy transport processes differ by region: radiation dominates in the core and radiative zone, where photons slowly diffuse outward. Convection takes over in the outer layers, rapidly moving energy to the surface. The photosphere emits visible light, while the chromosphere and corona involve complex wave and magnetic processes, which are vital for understanding phenomena like solar flares, sunspots, and the solar wind.

Energy Balance in the Photosphere: The Sun's photosphere energy balance links core energy production with the sunlight emitted. Energy generated by nuclear fusion in the core travels outward and is emitted as electromagnetic radiation at the photosphere. This balance helps explain the Sun's internal energy generation and the coronal heating paradox.

Energy Generation in the Core (Hydrogen Fusion): The Sun's energy comes from nuclear fusion in its core, where hydrogen nuclei fuse to form helium, releasing energy in the form of gamma rays and neutrinos. This energy is transported outward, eventually emitted as sunlight. The total luminosity of the Sun corresponds to the energy generated by fusion. The energy generation rate per unit mass describes the energy produced per kilogram of the Sun's core per second. The core's luminosity: $L_{core} = \varepsilon * m_{core}$, with ε as the energy generation rate (W/kg) and m_{core} as the core mass (kg), can be calculated and estimated. For the Sun, $\varepsilon \approx 2.5 \times 10^4$ **W/kg**, this value sets the baseline for the energy reaching the photosphere, though it takes thousands to millions of years for energy to diffuse outward.

Steady-State Energy Balance: In a steady state, the energy radiated from the photosphere matches the energy generated in the core. The luminosity is given by: $L = 4\pi R^2 \sigma T_{ph}^4$, where **L** is the luminosity, **R** is the Sun's radius, σ is the Stefan-Boltzmann constant, and T_{ph} is the photosphere's temperature. This equation shows that luminosity depends on the surface area of the photosphere and its temperature. The photosphere radiates all energy generated by fusion and even by energetic processes and forces we don't know or still not understand.

8. Heat Transfer in the Photosphere

The heat transfer in the photosphere is a delicate balance between radiation and convection. While radiation dominates the outward transfer of energy from the core, the convective zone just beneath the photosphere plays a significant role in shaping the Sun's surface structure and appearance. The temperature gradient across these layers is crucial for understanding how energy moves through the Sun's outer layers and contributes to phenomena like the solar granulation observed at the photosphere.

Convective Heat Transfer (Adiabatic Process): In the convective zone just beneath the photosphere, heat is transported by **convection**. The heat flux due to convection can be approximated by the following formula (for a convective layer): $Q = \rho * c_p * v * \Delta T$, where **Q** is the heat flux (W/m²), ρ the density of the plasma (kg/m³), c_p the specific heat capacity of the plasma (J/kg·K), **v** the convective velocity (m/s) and ΔT is the temperature difference between the convective material and the surrounding area (K). In the photosphere, the convective heat transfer is responsible for the appearance of granulation patterns, where hot gas rises to the surface and cooler material sinks. The **convective process** is responsible for the **granulation patterns** visible on the Sun's surface, where hot gas rises to the surface, loses heat, and sinks. These patterns are a direct result of the convective cells at the photosphere. The granules have

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

characteristic sizes of a few thousand kilometers, and their motions indicate a rapid heat transfer from the deeper layers to the surface. Convective heat transfer affects mostly the Sun's surface layers. Beyond the photosphere the behavior of energy becomes less straightforward, especially in the corona where **radiative transfer** and **magnetic interactions** dominate.

Temperature Gradient and Convective Instability: The temperature gradient in the convective zone is steep, with a drop of approximately 2 million K in the outermost part of the Sun's interior to around 5,500 K at the photosphere. The convective instability in this region is driven by the **Boussinesq approximation** to hydrodynamics, where the buoyancy force is the driving factor behind convection. The resulting convective cells transport heat more efficiently than pure radiation alone could, but the efficiency of this process decreases as the material becomes more transparent to radiation in the outer layers. Despite the efficiency of convection in the convective zone, the mechanism that transfers heat from the photosphere into the corona remains a central challenge.

9. Kinetic Theory of Plasma and Particle Interactions

The kinetic theory of plasma provides a more detailed, microscopic understanding of the behavior of plasma compared to fluid models. In the context of coronal heating, the kinetic theory is vital for explaining the fine-scale dynamics of particle interactions, energy transport, and dissipation processes in the corona. Fluid models often assume a continuum approach, where the plasma is treated as a smooth fluid, but this overlooks the discrete interactions between individual particles. The kinetic theory, on the other hand, explicitly accounts for these interactions and is essential for understanding phenomena like wave-particle interactions, collisions, and particle acceleration in the corona.

Boltzmann Transport Equation and Distribution Function: The key equation that governs the dynamics of particle distribution in a plasma is the **Boltzmann transport equation**. This equation describes how the distribution function of particles **f(v, r, t)** evolves over time under the influence of external forces and internal collisions. The equation is given by: $\partial f / \partial t + v * \nabla f + (F / m) * \nabla_v f = C[f]$, where **f(v, r, t)** is the **distribution function**, which gives the number density of particles at a given velocity **v**, position **r**, and time **t.** ∇f is the gradient of the distribution function in real space (m^{-4}) and **(F / m)** represents the force per unit mass (N/kg), with **F** as external force (e.g., electromagnetic force). $\nabla_v f$ is the gradient in velocity space, representing how the distribution of particles changes with respect to their velocities. **C[f]** is the **collision term**, which represents the rate of change of the distribution function due to collisions between particles or interactions between particles and waves. The collision term is crucial in the context of the coronal heating paradox because it governs the rate of energy dissipation in the plasma. In the corona, the collision frequency is generally low, but **wave-particle interactions** (through Alfvén waves, ion-cyclotron waves, etc.) can significantly affect the energy distribution, leading to **particle heating** and **wave damping**. These processes contribute to the overall **coronal heating** mechanism.

10. Magnetic Reconnection and Energy Release

Magnetic reconnection is a key mechanism in heating the solar corona. During reconnection events, oppositely directed magnetic fields reconnect, releasing energy in the form of thermal and kinetic energy, and accelerated particles. The **Sweet-Parker model** estimates the reconnection speed in a collisional plasma: $v_A = v_{ideal} * (L / d)$, where v_{ideal} is the Alfvén velocity, **L** the reconnection region's scale, **d** the current sheet thickness, and **ρ** is the plasma density. The velocity $(v_A = B / \sqrt{(\mu_0 \rho)})$ relates to the magnetic field strength and plasma density. Reconnection in the corona results in high temperatures, contributing to phenomena like solar flares. However, the Sweet-Parker model underestimates reconnection speed in high-energy environments, prompting the development of alternative models.

Modification with Lundquist number S: Faster reconnection at higher Lundquist numbers is a potential solution to coronal heating, with the dissipated magnetic energy heating the plasma. Numerical simulations, incorporating multi-fluid and kinetic effects, help improve understanding of these complex processes.

Observations of Solar Flares: Data from instruments like **GOES** and **SDO** show rapid energy release during solar flares, consistent with reconnection models. These processes correlate with specific X-ray and EUV emissions, providing direct evidence of heating.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

Petschek Reconnection Model: Unlike the **Sweet-Parker model**, the **Petschek model** predicts faster reconnection due to shock-like structures forming at the ends of the current sheet. The reconnection speed in the Petschek model is: **V_rec** $\propto$ **V_A / ln(L / δ)** , where **δ** is the ion diffusion region, and **L** is the reconnection region length. This model suggests faster reconnection speeds and more efficient energy release, enhancing coronal heating. Stronger magnetic fields and lower plasma densities accelerate reconnection.

11. Nanoflares and Magnetic Reconnection

Nanoflares are small bursts of energy that heat plasma in the corona to millions of Kelvin. They often coincide with high-energy electron populations, which are accelerated during magnetic reconnection. These electrons deposit their energy into the surrounding plasma, contributing to coronal heating.

Energy Dissipation during Reconnection: Energy released in magnetic reconnection is mainly thermal, heating the plasma. The total power dissipated during reconnection is: **P_reconnect = η J^2 V** , where **η** is plasma resistivity, **J** is current density, and **V** is the reconnection region volume. This dissipation is crucial in solar flares and other active regions, driving coronal heating.

Energy Release during Magnetic Reconnection: Magnetic reconnection occurs at various scales, releasing energy through heat and high-energy particles. The energy released is proportional to current density, magnetic field strength, and reconnection speed: **E_reconnect = η * (J * B) * V** , where **η** is the resistivity of the plasma, **J** the current density, **B** the magnetic field strength and **V** is the reconnection velocity. Magnetic reconnection remains one of the key drivers of coronal heating. This process leads to the formation of **hot plasma regions** in which ions are rapidly heated. During **solar flares**, magnetic reconnection releases vast amounts of energy, accelerating particles and ionizing species like **Fe^{6+}** and **O^{6+}** to high temperatures. These energetic particles are trapped in **magnetic loops**, where they interact with the plasma, causing further ionization and contributing to radiative cooling. Magnetic reconnection also plays a major role in the acceleration of **Coronal Mass Ejections (CMEs)**, which are massive bursts of solar wind carrying charged particles from the Sun's corona. These ejections can lead to **enhanced ionization** in the corona as the released energy heats the surrounding plasma. The amount of energy released during these processes can be quantified using models of **magnetic flux**, such as the **Sweet-Parker model** or the **Petschek model** of reconnection. The energy conversion efficiency is directly related to the ionization state of the plasma and the density of charged particles like Fe^{6+} and O^{6+}. Read more about solar flares and CMEs in the Sun's Water study and research papers for solar science.

Nanoflare Statistics and Energy Contribution: Although individual nanoflares release small amounts of energy, their cumulative effect can significantly contribute to maintaining coronal temperatures. Statistical analyses and numerical simulations suggest that nanoflares, occurring on a sub-minute timescale, could release energies ranging from **10^{8} to 10^{13} erg**, enough to sustain the corona's high temperature. Data from instruments like Hinode and the Parker Solar Probe support this idea.

Non-Thermal Electron Heating: Non-thermal electrons, accelerated by magnetic reconnection or wave-particle interactions, play a major role in coronal heating. These electrons transfer energy to the plasma via collisions and wave-particle interactions, contributing to the coronal heating paradox by raising the corona's temperature without direct heating from the photosphere.

Electron Energy Distribution: The energy distribution of accelerated electrons follows a power-law: **f(E)** $\propto$ **E^−α** , where **α** typically ranges from 1.5 to 2.0 for flare-associated electrons. The rate of energy deposition depends on electron density and collision rates with the ambient plasma: **Pe = ne * σe * ve * ΔE** , where **ne** is the electron density, **σe** the electron collision cross-section, **ve** the electron velocity and **ΔE** is the energy deposited in the plasma.Nanoflares, with their small-scale energy bursts, contribute to coronal heating through electron acceleration and energy deposition.

Shock Acceleration: In solar flares and CMEs, shock waves accelerate particles to high energies. These shocks, driven by reconnection and other dynamic processes, lead to a power-law distribution of particle energies: **f(E) = f_0 * (E / E_0)^(−α)** , where **E** is the energy, **f_0** is a normalization constant, and **α** is the power-law index.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

Stochastic Heating: In this process, electrons gain energy from turbulent magnetic fluctuations caused by reconnection. The Kolmogorov turbulence model describes how energy cascades from large to small scales, heating particles in the process. The energy distribution of electrons in a turbulent field is given by: **f(E) ∝ E^(-γ)** , where **γ** depends on turbulence level.

12. Non-Linear MHD Turbulence and Wave Dissipation

Magnetohydrodynamic (MHD) turbulence is central to the theory of coronal heating. It involves chaotic, non-linear interactions between magnetic fields and plasma, leading to the dissipation of wave energy as heat. MHD turbulence follows a power-law distribution, where energy cascades from large to small scales, eventually dissipating as heat. The energy density at a wavenumber **k** is: **E(k) = C * k^(-5/3)** , where **C** is a constant. This turbulence, observed in both the solar wind and the corona, is crucial for heating the plasma.

Dissipation of Alfvén Waves in the Corona: Alfvén waves, which propagate from the solar surface into the corona, dissipate their energy through non-linear wave damping or resonant absorption. This process is influenced by plasma density and magnetic field configuration. Resonant absorption occurs when waves become resonant with plasma at specific heights, transferring energy from the wave to the plasma.

Wave Damping by Resonance: The rate of wave energy dissipation in a resonant plasma is given by: **γ = (ω² * δ) / (B²)** , where **γ** is the damping rate, **ω** is the wave frequency, **δ** is a factor related to plasma density and magnetic field strength, and **B** is the magnetic field strength. This dissipation increases plasma temperature by converting wave energy into thermal energy.

Observational Evidence: Turbulence leads to current sheets and high-frequency waves, which dissipate energy as heat. Data from **SOHO** and **SDO** show significant turbulence in the corona, with wave-breaking and current sheet formation, supporting the idea that turbulence is essential for energy transfer and dissipation in the solar corona.

13. Observational Evidence and Empirical Models

Empirical data from solar missions, including **Hinode**, **IRIS**, **SDO**, and **SOHO**, have clarified the role of various heating mechanisms in the corona. More details on these missions are provided in the following sections. The integration of key findings, coronal seismology, high-resolution imaging, and spectroscopic data has enabled the refinement of theoretical models of coronal heating. Observations of solar flares, magnetic reconnection, wave propagation, and particle acceleration provide direct evidence that these processes contribute to corona heating. **High-resolution SDO data** have been instrumental in advancing our understanding of coronal heating. The **AIA** and **EVE** instruments aboard **SDO** have captured detailed images of solar flare emissions and the magnetic fields involved. These observations confirm that energy is released and dissipated in the corona through magnetic reconnection, wave-particle interactions, and Alfvén wave dissipation. Similarly, **Hinode** has provided crucial data on coronal loops, revealing oscillations and magnetic reconnection events contributing to heating. **IRIS** has delivered spectral data showing energy flow through the transition region between the chromosphere and corona. These insights and observations highlight fast magnetic reconnection events and small-scale plasma heating, reinforcing the idea that local reconnection significantly contributes to coronal heating. In addition to direct observations, numerical simulations based on empirical data have further validated these heating mechanisms. Models simulating wave propagation, magnetic reconnection, and turbulence replicate the observed temperature distribution in the corona, supporting the mechanisms maintaining high temperatures. Detailed research data can be obtained through **ESA and NASA's** official communication channels.
Empirical observations from solar missions have provided critical insights into spatial and temporal variations in the Sun's corona, supporting theoretical models of coronal heating. This section summarizes key findings and empirical evidence. These scientific approaches offer solutions to understanding coronal heating and other physical processes. By integrating research papers, references, studies, and key data, many challenges can be addressed. This article and advanced research from the solar science study contribute to developing a unified coronal heating theory, summarizing key mechanisms in the solar atmosphere and photosphere. The author works on a book called **"The Corona Paradox"**, this article

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

contains some parts of the first chapers. The scientific project and work will improve our understanding of the Sun's inner layers and the heliosphere, particularly regarding sunlight, solar winds, radiation, and particles. These advances will accelerate progress in planetary science, solar research, and space exploration.

Chromospheric and Coronal Coupling in Active Regions: Recent observations from the **Atacama Large Millimeter Array (ALMA)** and **Interface Region Imaging Spectrograph (IRIS)** have provided invaluable data linking the chromosphere to the corona. These observations suggest that energy transfer between the chromosphere and corona is critical in resolving the coronal heating paradox. **ALMA data** show that magnetic fields in active regions are highly dynamic, enabling continuous energy transfer between these layers. Energy is transferred through field-aligned currents, which assist in heat conduction upward, leading to localized heating and potentially micro-flares or small-scale magnetic reconnection events in the corona. **IRIS** has captured spectral data on chromospheric jets in active regions, showing that these jets channel energy into the corona via magnetic reconnection. The energy transferred from the chromosphere is significant enough to contribute to coronal heating when combined with wave and particle interactions. Chromospheric coupling, therefore, plays a key role in heating the corona. This energy exchange via magnetic reconnection and dynamic plasma flows is essential for maintaining the high temperatures of the solar corona.

Coronal Bright Points and Small-Scale Heating Events: High-resolution imaging of the solar corona has revealed **coronal bright points (CBPs)**, which are small, localized regions of enhanced brightness linked to transient heating events. Observations from **Hinode's EUV Imaging Spectrometer (EIS)** and **SDO's Atmospheric Imaging Assembly (AIA)** have identified intense heating in these areas, often associated with small-scale magnetic flux concentrations in the solar atmosphere. These bright points are thought to result from localized magnetic reconnection and current sheet formations, contributing to small-scale or transient heating events. These observations help refine our understanding of how the corona reaches its high temperatures without large-scale or global heating processes. Coronal bright points are primarily driven by localized magnetic reconnection, where oppositely directed magnetic fields release stored energy as heat. Regions with high electrical current density (current sheets) formed during reconnection are a source of Joule heating, contributing to localized temperature increases. Correlations between CBPs and magnetic flux concentrations suggest that small-scale processes, like magnetic reconnection and current sheet formation, are critical in heating the corona. These findings challenge traditional models based on large-scale heating, supporting the idea that localized processes, such as micro-flares and reconnection events, are essential to coronal heating. Observing these small-scale phenomena in high resolution enhances our understanding of fine-scale interactions in the solar atmosphere and their role in the corona's energy balance.

Coronal Loop Dynamics: Observations of coronal loops reveal complex magnetic structures that likely contribute to localized heating through magnetic reconnection. These loops, often exhibiting temperatures of 1–2 million K, support the hypothesis that reconnection is responsible for heating in these regions. Coronal loops act as energy storage reservoirs, accumulating energy that is later released via magnetic reconnection or wave heating. Observations from the **Parker Solar Probe** and **SDO** show that coronal loops can trap hot plasma, storing energy before releasing it. AIA observations have revealed that these loops undergo dramatic temperature fluctuations, with localized heating spikes near their footpoints. The loops focus wave energy from the lower atmosphere, concentrating it at higher altitudes. These oscillations and reconnection events in coronal loops contribute to solving the coronal heating paradox by storing and releasing energy in a controlled manner.

Coronal Mass Ejections (CMEs) and Solar Flares: Solar missions like **Hinode**, **SDO**, and **SOHO** have provided extensive data on solar flares and CMEs. **SOHO's LASCO** instrument has captured detailed images of CMEs, while **SDO's AIA** has monitored the solar corona during flare events. These observations reveal dramatic changes in the coronal magnetic field, consistent with the theory that magnetic reconnection releases energy during these events. **SDO's EVE** has detected large X-ray and ultraviolet emissions from solar flares, directly indicating energy release during magnetic reconnection. High-energy particles accelerated during flares are detected by **instruments like ACE**, providing evidence of particle acceleration and corona heating. Hinode's X-ray observations of flare events show localized heating in the corona, with temperatures reaching millions of degrees. These findings confirm that magnetic reconnection releases large amounts of energy into the corona, contributing to both explosive events like CMEs and the steady heating that sustains the corona's high temperatures.

Coronal Stratification and Magnetic Field Dynamics: The corona's temperature stratification is influenced by complex interactions between magnetic fields, wave dissipation, and non-radiative heating mechanisms. Energy is transferred from the chromosphere to the higher layers of the corona, where magnetic fields become more dynamic. In these regions, energy is deposited through wave dissipation, magnetic reconnection, and turbulence. Recent observations from **SDO** and **SOHO** have provided insights into coronal stratification, showing that the corona's density decreases rapidly with height, consistent with the observed temperature gradient. **SDO's AIA** has revealed signs of magnetic reconnection and turbulence in the lower corona, contributing to plasma heating. **Hinode's spectral data** show temperature and density variations across the corona, with observed coronal loops following a pattern where temperature increases with height in areas of strong magnetic activity. These observations support the idea that wave propagation and dissipation processes are responsible for maintaining the corona's stratified structure. High-resolution data from **Hinode's X-ray Telescope (XRT)** and **SDO** suggest that magnetic fields play a crucial role in sustaining the observed stratification, channelling energy from reconnection events into localized high-temperature regions. The **Global Oscillation Network Group (GONG)** provides complementary ground-based data on solar magnetic activity, supporting these findings.

Magnetic Reconnection and Energy Transfer: Direct empirical evidence for magnetic reconnection in the corona comes from observations of solar flares and CMEs. These explosive events are associated with rapid energy release, indicating that magnetic fields play a key role in coronal heating. Instruments like Hinode and SDO have captured detailed images of magnetic structures in the corona, showing dynamic magnetic loops and fields that facilitate energy transfer. **Hinode's XRT** has provided high-resolution X-ray images of reconnection jets, showing the release of energy that heats the plasma to millions of degrees. Observations from SDO reveal the rapid reconfiguration of magnetic field lines during solar flares, indicating magnetic reconnection. **SOHO's LASCO** has observed CMEs linked to magnetic reconnection events in the lower corona and upper photosphere layers. Radio emissions from these events support the idea that reconnection accelerates particles, contributing to corona heating. These observations align with models suggesting that magnetic reconnection drives both localized heating and global energy transport in the corona. Recent numerical simulations based on observational data have provided further insights, indicating that reconnection rates vary depending on the magnetic field structure and turbulence, explaining why heating is uneven across the corona.

More Evidence and Observations: Empirical evidence supporting the role of nanoflares and magnetic reconnection in coronal heating has emerged from several space-based instruments and missions, including the **Solar and Heliospheric Observatory (SOHO)**, **TRACE (Transition Region and Coronal Explorer)**, and more recently, the **Parker Solar Probe**. These instruments have detected small-scale energy bursts that exhibit characteristics consistent with magnetic reconnection and nanoflare activity in the Sun's corona. One key piece of evidence comes from the detection of high-frequency oscillations and rapid changes in coronal heating that coincide with reconnection events. Instruments such as **Extreme Ultraviolet Imaging Telescope (SOHO's EIT)** and **TRACE** have captured high-resolution images and spectroscopic data that reveal the dynamics of the corona in unprecedented detail. These observations show the presence of small-scale reconnection events, which are capable of releasing sufficient energy to heat the plasma to several million Kelvin, consistent with the temperature of the corona. As previously discussed, **magnetic reconnection**, **wave heating**, **wave-particle interactions**, and **turbulence** are the primary contenders for explaining the high temperatures observed in the corona. These mechanisms are capable of transporting energy from the Sun's photosphere to the corona, where it is dissipated as heat, overcoming the inefficiencies of traditional conductive heating models. The observational evidence supporting these mechanisms, including high-resolution images, spectroscopic data, and **solar wind measurements**, supports the conclusion that **non-radiative processes** (such as reconnection and wave heating) are primarily responsible for maintaining the elevated temperatures in the corona.

The Solar Cycle and Coronal Heating: The **11-year solar cycle** of solar activity has long been known to influence the intensity and frequency of **CMEs**, **solar flares**, and **sunspots**, and recent studies have explored how this cycle affects the long-term heating of the corona. During periods of **solar maximum**, when solar activity peaks, coronal heating is significantly enhanced. The frequency of solar flares and the intensity of magnetic reconnection events in the corona increase during these periods, leading to greater heat dissipation and higher coronal temperatures. **Parker Solar Probe** observations have revealed a direct correlation between flare frequency and coronal temperature during solar maximum, suggesting that these events are a major driver of energy input into the corona.

Conversely, during the **solar minimum**, when solar activity is subdued, heating in the corona appears to be more diffuse, primarily driven by turbulent and wave-related heating mechanisms. Data from the **SDO** and **Solar Orbiter** show that, even though solar activity is lower, the corona maintains elevated temperatures through the persistence of small-scale heating processes like nanoflares and Alfvén waves. These non-flaring heating mechanisms remain crucial for sustaining coronal temperatures during quieter phases of the solar cycle.

Numerical Simulations and Wien's Law: Numerical simulations based on observational data have offered further insights into coronal heating mechanisms. For example, the application of **Wien's Law** to the observed spectrum of the Sun has been used to study the coronal heating paradox. By comparing the observed spectral distributions from the photosphere, chromosphere, and corona, researchers have concluded that the energy output in the corona, especially in the UV and X-ray ranges, cannot be explained by traditional conductive heating models. Instruments like the **Hinode EUV Imaging Spectrometer (EIS)** have directly measured the spectral lines in the corona, supporting the idea that high-energy processes such as **wave dissipation** and **magnetic reconnection** are responsible for the observed temperature differences in the corona. High-resolution imaging from instruments like **DKIST (Daniel K. Inouye Solar Telescope)** and **SDO** has provided new insights into the role of small-scale magnetic structures, such as **magnetic flux sheets** and **magnetic reconnection events**, at the smallest scales of the corona. **DKIST** observations, in particular, have shown that small-scale magnetic structures (on the order of tens of kilometers) undergo rapid magnetic reconnection events. These events are responsible for both localized heating and the generation of **coronal jets**, suggesting that **micro-scale events** play a significant role in coronal heating. The dissipation of energy through these small-scale reconnection events has been found to be much larger than previously expected, providing new evidence that small-scale magnetic dynamics and wave-particle interactions contribute significantly to the heating of the corona.

Nanoflares and Small-Scale Events: High-resolution imaging from **SDO** has provided compelling evidence for **nanoflares.** These are small, localized bursts of energy that are hypothesized to be the primary mechanism for the continuous heating of the corona. The cumulative effect of numerous **nanoflares** and **microflares** could account for the observed high temperatures in the corona. Spectroscopic observations from **IRIS** and **SDO** show emissions consistent with the occurrence of nanoflares, particularly in regions of the corona with intense magnetic field activity, such as around **active regions** and **sunspots**. These regions exhibit frequent magnetic reconnection processes, supporting the theory that **small-scale magnetic reconnection events** are crucial for sustaining the high temperatures of the corona. Detections of nanoflares are not just theoretical, it has been confirmed as a real phenomenon. Observations show that these small-scale reconnection events lead to significant local heating, raising the temperature of the plasma to several million degrees Kelvin. This localized heating, resulting from magnetic reconnection events, contributes to the overall heating of the corona and may provide a solution to the **coronal heating paradox** by explaining the continuous energy input needed to maintain the high temperatures observed.

Scaling Laws and Observational Evidence: Empirical observations from **Hinode**, **SDO**, and **SOHO** offer direct evidence for many of the processes involved in coronal heating. By analyzing **scaling laws** derived from these observations, scientists have developed models to further support theoretical mechanisms of coronal heating. One such scaling law suggests that **magnetic field strength** and **plasma density** should be correlated with the observed **coronal temperature**. The empirical scaling law for the coronal temperature, with respect to magnetic field strength and plasma density, can be expressed as:
$T_{corona} = (B^2 * n)^{\alpha}$, where α is a constant derived from observational data. B is the magnetic field strength and n is the plasma density. This scaling law indicates that regions with stronger magnetic fields and lower plasma density, such as active regions and coronal holes, should exhibit higher temperatures, which is consistent with observations. **Coronal holes**, which are areas of the Sun's atmosphere with lower-density plasma and weaker magnetic fields, have been shown to maintain elevated temperatures, contrary to expectations based on their sparse magnetic activity. Recent observations from **Solar Orbiter** have revealed that the outflows from these coronal holes, which contribute to the **solar wind**, are thermally enhanced through local heating mechanisms. Studies using radio imaging techniques have shown that even though these regions lack intense flare activity, they still exhibit persistent low-level heating, driven by small-scale reconnection and wave-induced processes.

Solar Flares and Large-Scale Events: During solar flares, the release of magnetic energy leads to a sharp increase in the flux of non-thermal electrons, which can be directly observed through **hard X-ray emissions** captured by high-tech instruments like the **Ramaty High-Energy Solar Spectroscopic Imager (RHESSI)**.

These observations confirm that non-thermal electrons play a significant role in heating the plasma during flare events. **SDO's AIA** and **RHESSI** have captured detailed imaging and spectroscopy of flares, revealing how the release of magnetic energy accelerates particles and heats plasma to tens of millions of degrees within minutes. The relationship between **solar flares**, **CMEs**, and coronal heating is also evident in space weather data. Observations from space weather missions, such as **NOAA's DSCOVR** and **ACE**, have measured the impact of CMEs on Earth's magnetosphere, showing how energy is transferred from the corona into interplanetary space. These events provide crucial opportunities to test theories of coronal heating in real-time, as they involve large-scale energy release through **magnetic reconnection**.

Spectroscopic Observations: Spectroscopic data from **Hinode**, **SOHO**, and **SDO** provide key insights into the high temperatures in the corona. For example, spectral lines from highly ionized elements such as **Fe XII (iron)** and **Si VII (silicon)** provide direct measurements of the ionization state of plasma at different altitudes in the corona. These measurements confirm the high temperatures required for the proposed heating mechanisms, particularly in regions with strong magnetic fields. Coronal seismology, which analyzes the oscillations in coronal loops, has further supported the idea that **MHD waves** contribute significantly to coronal heating. **Wave energy dissipation** plays a major role in transferring energy from the photosphere to the corona, and **SDO's AIA** and **Hinode's EIS** have revealed that coronal loops experience **non-equilibrium heating** during flare events and magnetic reconnection processes. Spectroscopic data from **IRIS** have also shown elevated electron densities and temperature gradients in coronal loops, providing direct evidence that localized heating within these structures contributes to the overall heating of the corona.

Solar Radiation and Radiative Processes: Observational data from the **Hinode, SDO**, and **SOHO** have provided substantial evidence that the corona is heated through radiation and non-radiative mechanisms. The high-resolution imaging provided by these missions shows dynamic, complex magnetic structures that indicate the role of **magnetic fields** in the coronal heating process. Furthermore, **SOHO's** ultraviolet and X-ray observations reveal significant emission from the corona, which is consistent with very high-energy processes occurring in the corona, rather than simple radiative processes. Data from the **Hinode** satellite's **X-ray telescope** also support the idea that heating occurs in localized regions of the corona, where magnetic fields are intense and dynamic. **Cronal seismology** has provided evidence for the presence of waves which contribute to the heating processes. These waves can transport large amounts of energy from the photosphere and lower layers of the Sun's atmosphere to the corona, where they dissipate, contributing to the observed temperature rise. By analyzing wave patterns observed by instruments such as **Hinode** and **SDO**, scientists have confirmed the presence of **Alfvén waves** and **magnetoacoustic waves**, which are thought to be important contributors to the heating of the corona. These waves are observed to propagate from the photosphere upwards into the corona, with their energy being transferred through **wave-particle interactions** and **resonant absorption** mechanisms.

Supergranulation and Transition Regions: The role of supergranulation in coronal heating has become increasingly important in recent research. Supergranules, which are large convection cells observed on the Sun's surface, may influence the flow of energy from the photosphere to the corona, affecting the mechanisms responsible for coronal heating. **Hinode** and **SDO** have provided high-resolution images of **supergranular motions**, which reveal that these large convection cells could enhance the transport of energy to higher layers of the solar atmosphere, including the **transition region** and the **corona**. These motions may facilitate the generation of **Alfvén waves** and **magnetoacoustic waves** in the photosphere, which then propagate upward into the corona, carrying energy and contributing to the overall heating of the solar atmosphere. While the direct influence of supergranulation on the heating of the corona remains an active area of research, it is clear that the convective flows at the surface of the Sun play an important role in **generating the waves and turbulence that drive coronal heating**. The flow of energy from the photosphere to the corona through supergranular motions adds to the complexity of coronal heating mechanisms, contributing to the cumulative energy deposition that sustains the high temperatures observed in the corona.

The Role of the Transition Region in Coronal Heating: The transition region is the narrow layer between the Sun's **chromosphere** and **corona**. It has been found to play a crucial role in the process of energy transfer from the lower layers of the solar atmosphere to the corona. This region is characterized by rapid temperature changes, from about 10,000 K in the chromosphere to over a million degrees Kelvin in the corona. Understanding the dynamics of the transition region is essential for explaining how energy is transported across these layers, feeding into the heating of the corona. Recent observations from instruments like **IRIS** and **SDO** have revealed that the transition region undergoes significant heating events,

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

which contribute to the overall heating of the corona. Detailed spectroscopic data from these instruments show that plasma in the transition region is rapidly heated over very small spatial scales. This rapid temperature increase is consistent with **wave heating** and the interaction of plasma waves in the chromosphere, where energy is transferred upward to the corona. The transition region's unique structure presents a significant challenge for solar physicists. Due to its narrow spatial scale and sharp temperature gradients, studying this region requires extremely high-resolution instruments. **IRIS**, for example, is capable of capturing emission lines from the chromosphere and transition region, providing valuable data on the **energy transfer processes** that occur in these critical layers of the solar atmosphere. The sharp gradient in temperature from the chromosphere to the corona suggests that the transition region is a key zone for energy dissipation and conversion, where energy from the photosphere is rapidly transferred into the corona.

Some Future Directions and Observational Challenges: Despite the significant advances in our understanding of coronal heating mechanisms, many aspects of the process remain uncertain. Key questions remain about the relative contributions of different heating mechanisms, such as **nanoflares**, **wave heating**, **magnetic reconnection**, and **turbulence**, and how these mechanisms interact on both small and large scales to maintain the high temperatures of the corona. Future missions and observations will be crucial for refining our models of coronal heating. The **Parker Solar Probe**, **Solar Orbiter**, and **IRIS** are already providing invaluable data, and upcoming instruments such as **Daniel K. Inouye Solar Telescope (DKIST)** will offer even higher-resolution imaging, allowing scientists to study the Sun's atmosphere with unprecedented detail. **Parker Solar Probe**, in particular, will provide a closer look at the Sun's outer atmosphere, providing direct measurements of the solar wind, magnetic fields, and heating processes in the corona. One major challenge moving forward is integrating the various observational data with numerical simulations to create a comprehensive model that can explain the long-term behavior of the corona's temperature and the energy transport processes that sustain it. By combining data from space-based observatories, ground-based telescopes, and new solar missions, scientists are poised to uncover more detailed insights into the coronal heating paradox and its resolution.

Turbulence and Energy Dissipation: High-resolution data from **Hinode** and **SDO** have revealed small-scale turbulence within coronal loops, linked to magnetic reconnection events. These turbulent motions, observed as oscillations and plasma movements, provide direct evidence of turbulence in the corona. Regions of intense turbulence are often associated with high-energy flares and localized heating. **Hinode's EIS** has shown signs of turbulence, with spectral line broadening as a signature. This broadening results from Doppler shifts caused by turbulent plasma motions, confirming that turbulence plays a significant role in energy dissipation in the corona. **Numerical simulations** based on these observations indicate that turbulence amplifies magnetic reconnection, increasing energy transfer from large-scale structures to smaller heated regions. Turbulence also contributes to the formation of small-scale current sheets where energy is dissipated and transferred to the plasma. These findings suggest turbulence is crucial in both short- and long-term coronal heating. **SDO**'s coronal seismology data show oscillations likely driven by turbulence, contributing to energy dissipation. This evidence supports the idea that turbulence not only results from energy processes in the corona but also helps dissipate energy, fueling the heating of the solar atmosphere. Further data from **Hinode's SOT** and **SDO** reveal turbulent features correlated with localized heating. Small-scale turbulence, including eddies and vortices in the corona and solar wind, contributes to energy dissipation and the heating process. **IRIS** data also show turbulent motions in the transition region between the chromosphere and corona, feeding energy into the corona. This highlights turbulence's role in the solar atmosphere's dynamic behavior and sustained coronal heating. Numerical simulations further suggest that small-scale turbulence can heat both electrons and ions, potentially explaining the high temperatures of the corona despite its low plasma density.

Wave Propagation and Energy Transport: Hinode and **SDO** provide key evidence of wave propagation through the corona. **Hinode's EIS** detected both **Alfvén** and **magnetoacoustic waves**, supporting the theory that these waves are crucial for energy transport to the corona. Doppler shifts observed in spectral lines indicate wave motion and energy transfer. Non-thermal broadening of emission lines suggests waves energize the plasma through resonant interactions. **SDO's AIA** captured high-resolution images of coronal loops, showing waves propagating along them, confirming their role in coronal heating. Studies using **solar seismology** techniques have confirmed that wave patterns in the corona match predictions from magnetohydrodynamic (MHD) models, proving waves can transport energy from lower layers to the corona. One significant observation from **SDO** is the detection of high-frequency oscillations in coronal loops, matching the frequencies of Alfvén waves. These waves likely dissipate energy via resonant absorption, transferring their energy to plasma particles as wave amplitudes increase. Additionally, **SOHO** detected

propagating disturbances in the corona, confirming that waves cause local heating through non-linear processes like wave breaking and current sheet formation. **IRIS** observed wave motion in the transition region between the chromosphere and the corona, showing that waves carry energy to the corona. **SOHO's CDS** also detected wave-like phenomena in the transition region, reinforcing the role of waves in energy transfer to the corona. **Wave heating models** are further supported by observations of **coronal mass ejections (CMEs)** from **SOHO** and **STEREO**, which show wave-like structures and energy transfer from lower layers to the corona and interplanetary space. **Numerical simulations** based on **SDO** observations demonstrate that wave propagation efficiently transports energy across the Sun's atmospheric layers, offering a potential solution to the coronal heating paradox. **Coronal seismology** from **Hinode** and **SDO** shows a relationship between wave energy and temperature distribution in the corona, providing empirical evidence of the energy transport mechanisms responsible for coronal heating. Instruments on **Parker Solar Probe**, such as **FIELDS** and **SWEAP**, also support the idea of wave-particle interactions contributing to coronal heating, with wave dissipation leading to plasma heating. Spectroscopic measurements of the solar wind by **IRIS** show that waves accelerate electrons, which may further contribute to the high temperatures of the corona.

14. Opacity and Energy Transport

Opacity plays a crucial role in determining how energy is transported through the Sun's outer layers, including the photosphere. In general, opacity is a measure of how easily radiation is absorbed and scattered by particles (like electrons, ions, or atoms) in the Sun's material. Data from **SOHO** and **SDO** show that high-energy radiation in the **UV** and **X-ray regions** is emitted from the corona, supporting the idea that wave heating mechanisms are responsible for maintaining the high temperatures. These observations are consistent with theoretical models that account for non-radiative heating processes.

Kramers' Law for Opacity (Approximation in Solar Interiors): For many materials in the Sun's outer layers, the opacity κ can be approximated by **Kramers' law**, which is an empirical formula that relates **opacity to temperature and density:** $\kappa = \kappa_0 * \rho * T^n$, **w**here κ_0 is a constant that depends on the material (m^2/kg), ρ is the density (kg/m^3), **T** the temperature (K) and **n** is an empirically determined constant (typically between 1.5 and 2). For hydrogen and helium in the Sun's photosphere, $\kappa_0 \approx 1.0 \times 10^{-2}$ m^2/kg and $n \approx 1.5$. This relationship shows that the opacity increases with both higher density and higher temperature. In the photosphere, opacity is high enough to prevent efficient radiation transfer, which is why energy has to move outward via convective transport. Opacity determines the effectiveness with which energy can escape from the Sun's interior. The opacity of the material in the photosphere is crucial for determining how radiation is transported through the Sun.

Photospheric Opacity and Electron Scattering: The opacity due to electron scattering can be approximated by: $\kappa es = 0.34A * \rho * T-3.5$, where κ_es is the opacity due to electron scattering (m^2/kg), **A** the atomic mass number (for hydrogen, $A \approx 1$), ρ the density (kg/m^3) and **T** is the temperature (K). The scattering becomes an important source of opacity in the Sun's outer layers, especially in the photosphere, where temperatures are lower compared to deeper regions. This opacity helps trap radiation from deeper layers and causes the energy to accumulate in the photosphere before escaping as visible light.

Rosseland Mean Opacity: The mean opacity is commonly used to describe the opacity in stellar interiors, including the photosphere. It is given by the following expression: $\kappa_R = (16\pi acT^3) / (3\kappa\rho \lambda^4)$, where κ_R is the Rosseland mean opacity (m^2/kg), **a** the radiation constant (5.67×10^{-8} $W/m^2 \cdot K^4$), **c** the speed of light (3.00×10^8 m/s), **T** the temperature (K), ρ the density (kg/m^3) and λ is the wavelength of radiation (m). In the photosphere, opacity depends on the temperature, density, and the specific properties of the materials present. The high opacity in the Sun's interior means that radiation is repeatedly absorbed and re-emitted by atoms, slowly diffusing out toward the photosphere.

15. Particle Acceleration and Energization

In addition to wave and magnetic field interactions, particle acceleration plays a crucial role in coronal heating. During magnetic reconnection events, particles such as electrons and protons are accelerated to high energies, contributing to the observed coronal temperatures. The reconnection is typically more efficient at current sheets where the magnetic field is highly concentrated. These sites often have very small

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

scales and are strongly localized. The dynamics of such **small-scale reconnection events** can lead to particle acceleration on timescales that are much shorter than the typical magnetic field evolution. The phenomenon of reconnection allows the Sun's magnetic field lines to reorganize and reconnect, often resulting in the **release of a tremendous amount of energy**. This energy is not only dissipated as heat in the corona but also accelerates particles like electrons and protons up to **relativistic speeds near the speed of light**. These **particles can contribute significantly** to the **overall temperature** of the corona, which is observed to be much hotter than the photosphere, despite being farther from the Sun's core.

Electron and Proton Heating: Electron beams accelerated during reconnection events are often the primary contributors to the observed high-energy component of the solar flare spectrum. These electrons can accelerate the ions through **Coulomb collisions** and energy transfer, heating the plasma to millions of Kelvin. Another primary mechanism of particle acceleration in the corona is **Fermi acceleration**, which occurs during magnetic reconnection, another process is the betatron acceleration. The total energy gained by an individual particle depends on its interaction with the magnetic field and the time it spends in the reconnection region. The result is an **energy distribution with a non-thermal tail**, contributing to the overall coronal temperature. In Fermi acceleration, particles undergo **repeated scattering by magnetic irregularities**, causing them to gain energy each time they collide with these irregularities. This process can lead to the **rapid acceleration of particles**, resulting in high-energy electrons and protons. The kinetic energy of a single particle, either an electron or proton, is related to its relativistic speed. **The energy of a single particle** can be described by the relativistic formula: $E_particle = \gamma * m * c^2 - m * c^2$, where γ is the Lorentz factor, **m** the particle mass and **c** is the speed of light. This energy can then be transferred to the surrounding plasma, further increasing the temperature of the corona. The **Lorentz factor** accounts for relativistic effects, which become significant at speeds close to the speed of light. When a particle is accelerated to relativistic speeds during reconnection, the kinetic energy it gains can be enormous. The high-energy particles transfer their energy to the surrounding plasma, further increasing the temperature of the corona. This process is thought to be a significant contributor to the coronal heating paradox, where the corona is much hotter than the underlying photosphere, despite being farther from the Sun's core.

Observational Evidence: HPC calculations and data of the **Parker Solar Probe** indicates that electrons contribute significantly to the solar wind acceleration, especially at high altitudes in the corona, where kinetic energy of the electrons is transferred into thermal energy through interaction with plasma waves. **SDO's EUV and X-ray imaging** of the solar corona reveals that electrons are directly heated during **wave-particle interactions**. High-energy electron distributions can show that electrons play a significant role in transferring energy to the surrounding plasma. The observed **non-thermal electron populations** in the corona, identified through **X-ray emissions**, suggest that these electrons are not only heated but also accelerated by wave-particle interactions. Observations of solar flares and CMEs provide also direct evidence of particle acceleration, as X-ray emissions and radio bursts are often observed during these events. These emissions are consistent with models of non-thermal electron populations generated through wave-particle interactions. Observations from **RHESSI** and **Fermi** spacecraft have helped identify the high-energy tails in electron distributions, reinforcing the connection between particle acceleration and coronal heating.

Thermalization of Accelerated Particles: After particles are accelerated to high energies, they eventually thermalize with the surrounding plasma. This process is crucial for transferring the energy of accelerated particles to the bulk plasma, **raising the overall temperature of the corona**. The thermalization time describes how quickly energetic particles lose their energy to the surrounding plasma and is an important factor in understanding how much energy from accelerated particles contributes to coronal heating. **The thermalization rate** can be described by the quasilinear diffusion theory, which describes the scattering of particles by electromagnetic waves: $\tau = (\gamma * m * c^2) / (n * \sigma * v)$, where τ is the thermalization time, γ the Lorentz factor of the particle, **m** the particle mass, **c** the speed of light, **n** the number density of the particles, σ the scattering cross-section and **v** is the particle velocity. Thermalization occurs rapidly for high-energy particles, transferring their energy to the surrounding plasma and raising the temperature in the corona.
The **collisional frequency** of energetic particles with ambient plasma is also crucial. In the corona, the mean free path of particles is long, reducing the collision rate. However, when the energy of particles becomes sufficiently high, collisions with background plasma result in energy deposition, which in turn heats the plasma. Read more about solar particle and solar wind acceleration in the Sun's Water Theory and study papers.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

16. Photosphere Temperature Gradient and Energy Transport

The temperature gradient in the Sun's outer layers is a fundamental factor in determining how energy is transported from the Sun's interior to the outer atmosphere. The behavior of energy transport changes significantly across the different layers of the Sun, which can be divided into the photosphere, convective zone, and radiative zone. Each layer plays a crucial role in the solar energy balance and contributes to our understanding of coronal heating. In the Sun's convective zone, energy is primarily transported by convection, where hot plasma rises toward the surface, and cooler plasma sinks back down. This creates convective currents that carry energy from the interior of the Sun toward the surface. **The temperature gradient in the convective zone can be approximated by: $\nabla T = (T_{surface} - T_{core}) / H_{conv}$** , where ∇T is the temperature gradient (K/m), **T_surface** is the temperature at the Sun's surface (around 5,800 K), **T_core** is the temperature at the core of the Sun (about 15 million K) and **H_conv** is the convective scale height, which is the characteristic distance over which the temperature changes significantly due to convection (m). The convective zone extends from the **photosphere** down to approximately **0.7 solar radii**, where energy transport becomes dominated by radiation rather than convection.

Adiabatic Temperature Gradient (Convective Zone): In the convective zone, where convective transport dominates, the temperature gradient is determined by the **adiabatic** process (no heat exchange with surroundings). The temperature gradient can be approximated using the **adiabatic lapse rate for an ideal gas: $dT / dz = - (g * T) / (c_p * H)$** , where **dT / dz** is the temperature gradient (K/m), **g** the gravitational acceleration (m/s²), **T** the temperature (K), **c_p** the specific heat capacity at constant pressure (J/kg·K) and **H** is the pressure scale height (m), which is the height over which pressure decreases by a factor of **e.** In the photosphere, the convection is effective in transporting energy from deeper layers up to the surface, where the temperature drops sharply.

Effective Temperature (Photosphere Temperature): The temperature of the photosphere is an effective temperature, meaning it corresponds to the temperature of an ideal blackbody that emits the same amount of energy per unit area. It is related to the Sun's total luminosity by the **Stefan-Boltzmann law: $T_{ph} = (L / (16\pi\sigma R^2))^{(1/4)}$** , where **T_ph** is the effective temperature of the photosphere (K), **L** the luminosity of the Sun (W), **R** the radius of the Sun (m) and **σ** is the Stefan-Boltzmann constant. For the Sun, the temperature of the photosphere is approximately **5,800 K**, which is significantly cooler than the temperature of the corona.

Energy Transport in the Photosphere: In the photosphere, the temperature gradient is much smaller, and the primary mode of energy transport is radiation. However, once energy reaches the photosphere, it escapes as visible light and other wavelengths of radiation, which is the sunlight that we observe. The photosphere is the layer at which the mean free path of photons becomes large enough for them to escape without being reabsorbed, marking the transition from opaque to transparent material. **The mean free path for photons in the photosphere can be given by: $\lambda_{mfp} = 1 / (\kappa * \rho)$** , where **κ** is the opacity (m²/kg) and **ρ** is the plasma density (kg/m³). The mean free path indicates the average distance a photon can travel before it is absorbed or scattered. In the photosphere, this path is short, which leads to multiple scattering events before a photon escapes.

Temperature Gradient in the Radiative Zone: Below the convective zone lies the radiative zone, where energy is transported outward through radiation. In the radiative zone, the temperature gradient can be described by the radiative diffusion equation. The temperature gradient is determined by the opacity of the plasma, which is strongly dependent on the density and temperature of the material. **The gradient can be written as: $dT / dr = - (3\kappa\rho L) / (16\pi acR^2T^3)$** , where **dT / dr** is the temperature gradient (K/m), **κ** the opacity of the material (m²/kg), **ρ** the density (kg/m³), **L** the Sun's luminosity (W), **a** the radiation constant with 5.67×10^{-8} W/m²·K⁴, **c** the speed of light (m/s), **R** the radius of the Sun (m) and **T** is the temperature (K). In the radiative zone, energy is transported outward very slowly due to the high opacity of the material. Photons undergo frequent scattering, which results in radiative diffusion, leading to a temperature gradient that decreases as energy moves outward toward the photosphere.

17. Planck's Law and the Solar Spectrum

The solar spectrum, which encompasses all wavelengths of electromagnetic radiation emitted by the Sun, is governed by Planck's Law. This law tells us how the intensity of radiation varies with wavelength at a given temperature: **$I(\lambda, T) = (2\pi hc^2) / (\lambda^5) * (1 / (\exp(hc / (\lambda kT)) - 1))$** , where **I(λ, T)** is the spectral radiance

(W/m²·m·sr), **λ** the wavelength (m), **h** is Planck's constant (6.626×10^{-34} J·s), **c** the speed of light with 3.00×10^8 m/s, **k** the Boltzmann constant (1.38×10^{-23} J/K) and **T** is the temperature of the object (K). This law applies to the Sun's photosphere, showing how emitted intensity varies with wavelength. The peak wavelength of solar radiation corresponds to the visible light range (about 500 nm), which is why the Sun appears yellowish-white. The temperature of the photosphere is approximately 5,778 K. **Planck's Law** helps describe the solar spectrum, which spans a wide range of wavelengths from ultraviolet (UV) to infrared (IR) and includes the visible light range, where the Sun's radiation peaks. Data from space missions such as **SDO** and **SOHO** provide empirical evidence of the solar spectrum, showing that **UV** and **X-ray emissions** primarily come from the corona. Spectral shifts observed from these instruments confirm that the corona is much hotter than the photosphere, supporting the role of non-radiative heating.

Peak Wavelength of the Solar Radiation: The peak wavelength at which the Sun emits most of its radiation can be determined by **Wien's Displacement Law**, which relates the temperature of a blackbody to the wavelength at which it emits the most radiation: $\lambda_{max} = b / T$, where λ_{max} is the peak wavelength (m), **b** the Wien's constant (2.898×10^{-3} m·K) and **T** is the temperature of the blackbody (K). For the Sun, with a temperature of 5,778 K, the peak wavelength is: $\lambda_{max} = 2.898 \times 10^{-3} / 5{,}778 \approx 5.02 \times 10^{-7}$ **m = 502 nm !** This wavelength lies in the visible light range of the electromagnetic spectrum, which is why the Sun appears white or yellowish. The Sun's radiation peaks near the green part of the visible spectrum with approx 500 nm, but the combination of emitted wavelengths across the visible range results in the Sun's characteristic white light.

The Solar Constant and Total Emission: The total energy emitted by the Sun, or its luminosity, can be derived from the total emission of radiation across its entire surface, using the **Stefan-Boltzmann Law** for blackbody radiation. This law relates the total radiated power per unit area to the temperature of the blackbody: $L_{Sun} = 4\pi R_{Sun}^2 \sigma T_{Sun}^4$, where **L_Sun** is the luminosity of the Sun (W), **R_Sun** the radius of the Sun (6.96×10^8 m), **σ** the Stefan-Boltzmann constant (5.67×10^{-8} W/m²·K⁴) und **T_Sun** is the temperature of the Sun's photosphere (5,778 K). The total solar luminosity is approximately 3.828×10^{26} **W**, representing the total energy the Sun emits in all directions per unit of time. The **solar constant** is the amount of solar energy received per unit area at the Earth's distance from the Sun. It can be calculated by dividing the Sun's luminosity by the surface area of a sphere with a radius equal to the Earth-Sun distance: $S_0 = L_{Sun} / (4\pi d_{Earth\text{-}Sun}^2)$, where **d_Earth-Sun** is the distance from the Sun to Earth (approximately 1 AU, or 1.496×10^{11} m). The solar constant is approximately **1361 W/m²**.

The Solar Spectrum is a continuous spectrum that spans a wide range of wavelengths. It can be divided into several regions, including **ultraviolet (UV)**. The Sun emits significant energy in the UV range, which is crucial for processes like photosynthesis and the impact of solar radiation on Earth's atmosphere. The Sun emits a broad spectrum of visible light, with peak intensity in the green-yellow part of the spectrum. This is why the Sun appears white from space but can appear yellowish from Earth due to atmospheric scattering. The Sun also emits significant radiation in the infrared, which, though invisible to the human eye, contributes to Earth's heating and is important for the global energy balance. The real solar spectrum also contains **absorption features**, known as **Fraunhofer lines**, which are produced when specific elements (e.g., hydrogen, sodium, calcium) in the Sun's atmosphere absorb radiation. These absorption lines provide important information about the chemical composition and structure of the solar atmosphere.

18. Plasma Processes and Plasmonic Structures

Coronal mass ejections (CMEs) are large expulsions of plasma and magnetic fields from the corona into space. These eruptions release magnetic energy, primarily from magnetic field line reconnection. The energy of a CME can reach 10^{16} joules, much larger than typical solar flares. The rapid energy release during a CME disturbs the corona, with accelerated particles heating the surrounding plasma. Understanding CME interactions with the solar wind and magnetic reconnection processes is essential for explaining their role in coronal heating. During a CME, magnetic reconnection intensifies, and shock waves accelerate particles, which then transfer their kinetic energy to the plasma via collisions, contributing to coronal heating. The energy transfer rate in these collisions is explained by the Coulomb collision rate (section 20). CME particles collide with surrounding plasma, dissipating energy and increasing the local plasma temperature, especially in regions where shock waves interact with the solar wind. During CME events, plasmonic structures form due to wave-particle interactions. These structures, including plasma waves and oscillations, influence energy distribution within the ejected plasma. Plasmonic effects enhance local

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

heating, converting electromagnetic wave energy into thermal energy. Langmuir waves, a type of plasma oscillation, are particularly important. The frequency of these waves is given by the **plasma frequency:** $\omega_p = \sqrt{(n_e * e^2 / (m_e * \varepsilon_0))}$, where n_e is the electron density, **e** the electron charge, m_e the electron mass, and ε_0 the permittivity of free space. These waves transfer energy to plasma particles, enhancing coronal heating, especially during CME events.

Solar Flares and Magnetic Reconnection: Solar flares are driven by magnetic reconnection. While CMEs involve large-scale plasma expulsions, solar flares rapidly release electromagnetic radiation (X-rays, UV) and energetic particles. The energy from solar flares also heats the surrounding plasma, contributing to coronal heating dynamics. The energy released during a solar flare is given by: $E_f = \eta * \Delta V * B^2$, where E_f is the flare energy, η is the reconnection efficiency, ΔV is the volume undergoing reconnection, and **B** is the magnetic field strength. This energy is transferred as electromagnetic radiation and through particle acceleration, increasing the plasma's thermal energy. More details about solar flares and CMEs can be found in chapter 10 of the Sun's Water Theory.

Magnetic Reconnection and Energy Release

Magnetic reconnection in the coronal plasma converts magnetic energy into kinetic and thermal energy. This explosive energy release accelerates particles and ejects plasma. During a CME, reconnection intensifies over large scales. The released energy is given by: $E = (B^2 * V * L^3) / \mu_0$, where **E** is the released energy, **B** the magnetic field strength, **V** the ejected plasma velocity, **L** the CME's characteristic length, and μ_0 is the permeability of free space. The magnetic field influences particle acceleration and shock wave formation. Shock waves accelerate ions and electrons, which then transfer their energy to the coronal plasma, contributing to heating. Particle acceleration can be described by the Fermi acceleration process.

Plasma Conduction and Energy Transport

Heat transport in the corona is governed by thermal conduction, which moves heat from hotter to cooler regions. The classical conduction equation is: $Q = -\kappa \nabla T$, where **Q** is the heat flux, κ is thermal conductivity, and ∇T is the temperature gradient. In the corona, thermal conductivity is influenced by magnetic fields, restricting conduction along the field lines. This anisotropic conduction must be modeled accurately for better energy transport predictions. The modified equation is: $Q = -\kappa_{perpendicular} \nabla T_{perpendicular} - \kappa_{parallel} \nabla T_{parallel}$, where $\kappa_{perpendicular}$ and $\kappa_{parallel}$ represent the perpendicular and parallel components of the conductivity tensor. In the corona, $\kappa_{parallel}$ is much larger than $\kappa_{perpendicular}$ due to the alignment of magnetic fields with plasma's thermal energy. The magnetic field creates anisotropic heat transport, with energy mainly moving along the magnetic field lines. Magnetic confinement and heat conduction lead to high-energy regions in the corona, releasing energy via reconnection, wave dissipation, and particle acceleration.

Plasma Expansion and the Solar Wind: The thermodynamics of the corona also involve plasma expansion into the solar wind. As plasma expands, its density decreases, and temperature rises due to adiabatic expansion. The relationship between temperature and density is: $T * n^{(\gamma - 1)} = \text{constant}$, where **T** is the temperature, **n** is particle number density, and γ is the adiabatic index (5/3 for mono-atomic gases). As plasma density decreases in the corona, temperature rises, conserving energy. This process contributes to the high-speed solar wind observed at the Sun's surface, though additional heating mechanisms, like wave dissipation or magnetic reconnection, are needed to explain the corona's millions of degrees.

Plasma Heating by Collisions and Conductivity: While waves and reconnection dominate coronal heating, collisional heating and thermal conduction are important, particularly in the lower corona. The Spitzer conductivity governs heat transport in a plasma and affects the corona's thermal structure. Heat flux is: $q = -\kappa * \nabla T$, where **q** is heat flux, κ is thermal conductivity, and ∇T is the temperature gradient. In the corona, heat flux is large near the base but decreases at higher altitudes due to lower density. Spitzer conductivity at the base is high, leading to significant heat transfer, but the decrease in heat flux with height suggests other mechanisms, like wave dissipation or magnetic reconnection, are crucial for maintaining high temperatures. **The Spitzer conductivity is given by:** $\kappa_S = (5.8 \times 10^{-6}) T^{(5/2)} / n \ln \Lambda$, where κ_S is Spitzer conductivity, **T** is plasma temperature, **n** is particle density, and $\ln \Lambda$ is the Coulomb logarithm, which depends on plasma interactions.

Electron Collisions and Temperature Regulation: In high-density regions of the corona, electron-electron and electron-ion collisions regulate energy and further heat the plasma. The energy transfer rate from electrons to ions is: $P = n_e * n_i * (kT_e)^2 * \ln(\Lambda)$, where **P** is the power transferred, n_e and n_i

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

are electron and ion densities, **T_e** is electron temperature, **k** is the Boltzmann constant, and **ln(Λ)** is the Coulomb logarithm. Electron-ion collisions play a key role in maintaining the high temperatures in the corona by facilitating energy transfer between particles, especially in denser regions.

Mathematical Model for Stochastic Heating: Plasma wave-particle interactions can be modeled using the Fokker-Planck equation: **∂f / ∂t = ∂ / ∂p (D_pp ∂f / ∂p) + ∂ / ∂p (C(p) f)** , where **f(p,t)** is the particle distribution function, **D_pp** is the diffusion coefficient, and **C(p)** represents particle collisions. Non-thermal electron distributions observed during solar flares support wave-particle interactions as a key heating mechanism in the corona. These accelerated particles transfer energy to the plasma, raising its temperature. While wave heating and magnetic reconnection dominate, collisional heating and thermal conductivity are important for understanding the lower corona, where collisions are more frequent.

Plasma Waves and Instabilities: Plasma waves, including whistler waves, ion-cyclotron waves, and fast magnetosonic waves, are crucial for energy transport through the corona. These waves accelerate electrons, cause resonant absorption, and generate instabilities that dissipate energy into the plasma. Turbulent motions lead to micro-instabilities that enhance energy dissipation. The firehose instability, for example, occurs when plasma pressure exceeds the magnetic field, causing anisotropies in particle velocity distributions. The energy dissipation rate due to plasma instabilities is: **P = γ * (n_e * k_B * T)** , where **γ** is the instability growth rate, **n_e** is electron density, **k_B** is the Boltzmann constant, and **T** is the temperature. These instabilities accelerate particles, enhance energy dissipation, and drive complex interactions in the solar corona. Some key plasma instabilities in the solar corona include:

Kelvin-Helmholtz Instability (KHI): Occurs at the boundary of different plasma layers with different velocities. In the context of the corona, this can happen at the interface between the solar wind and the coronal plasma, or within coronal loops. KHI can lead to the **mixing of plasma layers**, causing **turbulent motions** that accelerate ions such as **Fe^{6+}** and **O^{6+}** and increase the energy dissipation through collisionless heating.

Rayleigh-Taylor Instability (RTI): This instability occurs when a lighter plasma is above a heavier plasma (e.g., the photosphere or transition region). The RTI can enhance the **mixing** and **reconnection** processes, providing pathways for **electrons** and **ions** to be energized. This instability is particularly important during **solar flares** and **coronal mass ejections (CMEs)**, as it leads to the rapid transfer of energy to both ions and electrons.

Tearing Modes: Associated with magnetic reconnection events, tearing modes can break up magnetic field lines and cause localized heating of the plasma. The localized reconnection increases the energy dissipation in the corona, contributing to both the **heating of ions** (such as Fe^{6+} and Mg^{6+}) and the acceleration of particles. Tearing modes are especially relevant in the **solar flare and coronal mass ejection (CME)** environments. These plasma instabilities contribute significantly to the non-equilibrium behavior of the solar corona, leading to highly dynamic processes that enhance the overall energy transfer and ionization balance.

21. Solar Spectrum and Frequencies of Light

The **solar spectrum** plays a crucial role in understanding the behavior and heating of the Sun's outer atmosphere, particularly the **corona**. Spectral lines, especially in the **UV** and **X-ray** regions, serve as valuable diagnostic tools for studying the **temperature**, **density**, and **velocity** of the plasma in the corona. These spectral lines are typically emitted by highly ionized species such as **Fe IX**, **Fe XIII**, and **Ca XV**, and they help researchers map the thermal structure of the corona. The ionized species correspond to different temperature ranges, providing insight into the dynamic conditions of the coronal plasma. By analyzing the observed spectral data, researchers can infer key properties like temperature gradients, density variations, and plasma flows, which are essential for understanding energy transport mechanisms in the corona. The energy balance of the corona depends heavily on the way in which energy is transferred from the **photosphere** through **radiation**, **wave heating**, and **magnetic reconnection**. While the photosphere emits radiation primarily in the visible range, a significant portion of solar energy is radiated in the **ultraviolet (UV)** and **X-ray** wavelengths. These higher-energy photons are absorbed by the corona, where they contribute to the ionization of particles and accelerate them, playing a key role in the observed coronal heating.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

The UV and X-ray emissions from the Sun's atmosphere are responsible for ionizing the coronal plasma and driving dynamic processes such as **wave-particle interactions** and **magnetic field reconfigurations**, both of which are thought to contribute significantly to coronal heating. The **Plank function** and **blackbody radiation** theories help explain the spectrum of solar radiation, with the **UV** radiation acting as a primary driver of the energy processes that sustain the high temperature of the corona. In terms of solar radiation, the photosphere radiates a broad spectrum of energy, peaking in the visible range, but extending across the **electromagnetic spectrum** into the **infrared (IR)** and **ultraviolet (UV)** ranges. The UV radiation, in particular, plays an essential role in heating the plasma, contributing to high-energy processes that further ionize and accelerate particles in the corona. The combination of these radiation processes explains much of the heat transfer in the solar atmosphere.

22. Solar Wind Contribution to the Corona and Photosphere Dynamics

The solar wind is a stream of charged particles (mostly protons and electrons) that emanates from the outer layers of the Sun, including the corona. While the solar wind's primary role is to transport energy and matter away from the Sun, it also has a subtle contribution to the dynamics of the photosphere and the **coronal heating** process. The connection between the solar wind and coronal heating is primarily driven by the acceleration of solar wind particles in the high-temperature regions of the Sun's outer atmosphere.

Acceleration of Solar Wind and Key Drivers: The acceleration of the solar wind is closely tied to the temperature gradient between the corona and the photosphere. The corona, which reaches millions of degrees Kelvin, is much hotter than the **photosphere** (~5,800 K). This large temperature difference creates a pressure gradient, which drives particles outward, forming the solar wind. Although this pressure difference is a fundamental driver, additional energy is supplied by processes in the corona, including **magnetic reconnection** and **wave dissipation**. The Sun's magnetic field, which extends through the corona, plays a critical role in shaping the solar wind's properties. **Magnetic reconnection** events in the corona release vast amounts of energy, which is converted into **thermal energy**. This thermal energy drives the temperature increase in the corona and accelerates the solar wind particles. Additionally, **Alfvén waves**, which propagate along the magnetic field lines, transfer momentum to the plasma, helping accelerate the solar wind. These waves are generated by the turbulence and dynamics associated with magnetic field interactions in the corona.

Energy and Mass Transport: The solar wind not only carries mass but also energy away from the Sun. As it moves outward, the wind transports both thermal energy and momentum, which were initially provided by the Sun's corona. The energy dissipation processes within the corona, including magnetic reconnection and wave interactions, help maintain the high temperatures of the corona by continually feeding energy into the solar wind. Without the transport of energy through the solar wind, the corona's temperature would decrease over time, as energy would accumulate near the photosphere. Therefore, the solar wind plays an essential role in maintaining the energy balance of the Sun's outer layers.

Solar Wind Mass Loss: The mass loss due to the solar wind is governed by the **continuity equation**:
$\mathbf{dm / dt = 4\pi r^2 \rho v_s}$, where dm/dt is the mass loss rate due to the solar wind (kg/s), **r** is the radius of the Sun (m), **ρ** the mass density of the solar wind (kg/m³) and **v_s** is the speed of the solar wind at a given distance (m/s). The solar wind is composed of protons, electrons, and heavier ions, all of which have been heated to high temperatures in the corona. As they move outward from the Sun, their **kinetic energy** contributes to maintaining the elevated temperatures of the corona, and only a small fraction of the energy flows back to the photosphere.

Sunspot Temperature and Magnetic Field Influence: While the **corona** is exceptionally hot, sunspots are cooler regions of the photosphere due to the influence of strong magnetic fields that suppress convection. The temperature of a sunspot is generally lower than that of the surrounding photosphere, which is approximately **5,800 K**. Sunspot temperatures are typically about **1,000–1,500 K** cooler than the photosphere, with sunspot temperatures reaching around **4,000 K**. This temperature difference is due to the blocking effect of magnetic fields on the convective heat transfer in the Sun's interior.

The Role of Solar Wind and Outflow Dynamics: The solar wind, while often overlooked in discussions of coronal heating, plays an important role in regulating the energy dynamics of the Sun's outer atmosphere. The wind, composed mainly of **ionized particles**, carries both energy and mass outward, shaping

the interaction between the corona and the interstellar medium. This interaction leads to dynamic effects such as **shock waves** and **compression zones**, which can further dissipate energy and contribute to the heating of the solar corona. The outflow dynamics of the solar wind influence **large- and small-scale** eruptions of plasma. The acceleration of the solar wind and the associated **magnetic reconnection events** help drive these massive bursts of energy, contributing significantly to coronal heating. The solar wind's speed and the energy it carries depend on local heating mechanisms in the corona, especially in **polar coronal holes**, where the **solar wind** is typically faster and more efficiently accelerated.

Solar Wind Acceleration and Magnetic Field Dynamics: The acceleration of the solar wind is intricately tied to the **magnetic field** configuration in the corona. In areas with **open magnetic fields**, such as **coronal holes**, the solar wind is more easily accelerated outward. These regions exhibit lower plasma density and higher temperatures, allowing more efficient acceleration. The lower plasma density facilitates the faster outflow of particles, leading to **high-speed solar wind** with velocities reaching **700–800 km/s**. The acceleration of the solar wind is described by the equation: $\mathbf{d/dt\ (1/2\ mv^2) = -\ GMm\ /\ r^2}$, where **m** is the mass of a solar wind particle, **v** the velocity, **G** the gravitational constant, **M** the mass of the Sun and **r** is the distance from the Sun's center. The increase in kinetic energy as solar wind particles are accelerated is a result of **magnetic field interactions** and energy dissipation in the corona. These processes help maintain the high temperatures in the corona by continually feeding energy into the outward flow of the solar wind.

23. The Role of Gravity and Coronal Stratification

Gravity plays an indirect yet fundamental role in coronal heating by shaping the structure of the solar atmosphere, particularly influencing the density distribution and the formation of the solar wind. The stratification of the solar corona, driven by the Sun's gravity, means that the density of the plasma decreases exponentially with height. This stratification impacts the temperature distribution in the corona and helps determine how energy is transported and stored within the outer layers of the Sun.

Barometric Equation for Corona Stratification: The vertical distribution of plasma density in the corona is governed by the **barometric equation**, which can be written as: $\mathbf{n(r) = n_0\ exp(-m\ g\ r\ /\ k_B\ T)}$, where **n(r)** is the particle density at radius **r** (m^{-3}), $\mathbf{n_0}$ the particle density at the base of the corona (m^{-3}), **m** the particle mass (kg), **g** the gravitational acceleration (m/s^2), **T** the temperature (K) and **k_B** is the Boltzmann constant (J/K). This exponential decrease in density with height implies that as plasma rises into the corona, it undergoes adiabatic expansion, affecting its temperature. However, this alone cannot explain the extremely high temperatures of the corona. Additional heating mechanisms, like wave dissipation, are needed.

Gravitational Stratification of the Corona: Gravity's influence on the Sun's outer layers leads to the **stratification** of the corona, which refers to the decreasing plasma density with height above the photosphere. This stratification is governed by gravity, as it holds the plasma in place while allowing it to expand outward as the temperature rises. The plasma density in the corona decreases by a factor of 10^4 to 10^5 over a height of about 1-2 solar radii, and this behavior can be described by the following **barometric formula**: $\mathbf{\rho(r) = \rho_0\ *\ exp(-GM\ /\ kTr)}$, where **ρ(r)** is the density at a distance **r** from the Sun, $\mathbf{\rho_0}$ the density at the photosphere, **G** the gravitational constant, **M** the mass of the Sun, **k** the Boltzmann constant and **T** is the temperature at height **r**. This stratification implies that the plasma density in the corona is very low compared to the photosphere, but the temperature increases with height. However, the increase in temperature with height, known as **coronal heating**, remains a paradox since the photosphere is cooler than the corona. This unexplained temperature rise is a critical challenge in solar physics.

Gravity's Influence on the Photosphere and Lower Solar Atmosphere: Gravity also influences the photosphere and the transition to the corona. At the photosphere, gravity-induced compression stabilizes the layer, maintaining a temperature of around 5,800 K. However, as plasma rises from the photosphere toward the corona, magnetic fields become more dynamic and can reconnect, releasing energy through solar flares and coronal mass ejections (CMEs). Gravity contributes to how these fields break and reconnect, indirectly influencing the heating mechanisms responsible for the high temperatures of the corona.

Temperature Distribution and Gravitational Effects: In a gravitationally stratified atmosphere, the temperature usually decreases with height due to the adiabatic expansion of rising plasma. However, in the corona, the temperature increases dramatically with height, creating a temperature gradient that challenges traditional expectations. This rise in temperature is crucial to understanding coronal heating and requires additional processes beyond gravity, such as magnetic and wave heating.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

Gravitational Effects on Plasma Heating: Near the Sun's surface, the gravitational force compresses the plasma, but as plasma rises into the corona, the weakening of gravity allows for expansion, which causes cooling in the lower regions. The radial temperature gradient in the corona is influenced by gravitational forces and weak magnetic fields. The interaction between gravity and magnetic fields plays a role in energy transfer and sustaining coronal heating.

Gravitational Settling and Thermal Conduction: At lower altitudes, heavy ions (e.g., **Fe** and **O**) settle toward the Sun's surface, while lighter ions (e.g., **H** and **He**) tend to rise. This differential settling impacts both the composition of the solar atmosphere and the temperature distribution. In a gravitationally stratified plasma, heat is carried upward through **thermal conduction**, but the tenuous nature of the corona means **Spitzer conduction** alone cannot explain the high temperatures. Additional mechanisms, such as wave-particle interactions and magnetic heating, are needed to account for the temperature rise. Gravity influences how these processes interact with the plasma, aiding in the transport of heat.

24. The Sun's Atmosphere Structure and Boundaries

The photosphere, which marks the visible "surface" of the Sun, is the layer from which light escapes. However, it is not a solid surface, and it represents a transition between opaque and transparent regions. Below the photosphere, the Sun's material is opaque to radiation. This boundary, between the photosphere and the layers above it, is governed by the mean free path of photons, which refers to the average distance a photon can travel before being absorbed or scattered. The **mean free path** of photons in the photosphere can be expressed as: $\lambda_{mfp}=1\kappa\rho$, where λ_{mfp} is the mean free path of photons (m), κ the opacity (m^2/kg) and ρ is the density (kg/m^3). In the Sun's photosphere, the mean free path is quite short (a few millimeters to a few centimeters) due to the high density of the plasma. This short mean free path results in a highly **opaque** photosphere, where photons undergo multiple scatterings before escaping into space.

Boundary Conditions and Coronal Heating: The transition between the photosphere and the corona is not sharp. Instead, it involves a complex zone where energy, plasma, and magnetic fields interact. As energy travels outward from the Sun's core through the radiative and convective zones, much of it is absorbed and re-emitted in the photosphere. However, some of the energy is directed into the corona, where it is converted into heat. Understanding how energy reaches the corona, despite the scattering and absorption in the photosphere, is key to resolving the **coronal heating paradox**. This energy is likely delivered by processes such as **magnetic reconnection**, **wave propagation** (including **Alfvén waves**), and **turbulence**, which allow energy to bypass the inefficient radiative transport mechanisms in the photosphere.

The Role of the Transition Region and Chromosphere: The chromosphere lies between the photosphere and the corona and experiences rapid temperature and density changes. The temperature rises from around 6,000 K in the photosphere to about 20,000 K in the chromosphere, and then skyrockets to millions of degrees in the corona. Magnetic fields are crucial in this region, influencing energy transport and plasma dynamics. Reconnection events, wave heating, and turbulence play a key role in transferring energy to the corona.

25. Turbulence in the Corona

Turbulence in the solar corona is another significant factor contributing to coronal heating. The chaotic motion of plasma, which forms turbulent flows, enhances energy dissipation, leading to higher temperatures in the corona. Turbulent flows can be described using the **Navier-Stokes equation** for an incompressible fluid, modified for plasmas. The general form of the equation is: $\partial \mathbf{v} / \partial t + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\nabla p + \nu\nabla^2\mathbf{v} + (\mathbf{J} \times \mathbf{B}) / \rho$, where $\partial \mathbf{v}/\partial t$ represents the time derivative of the velocity field (acceleration), $(\mathbf{v} \cdot \nabla)\mathbf{v}$ the convective acceleration (advection term), ∇p is the gradient of pressure and ν is the kinematic viscosity - describing the diffusion of momentum). $\nabla^2\mathbf{v}$ is the **Laplacian** of the velocity (describing the diffusion of velocity), $(J \times B)$ is the **Lorentz force** (interaction of the current density **J** and magnetic field **B**), and ρ is the plasma density. This equation governs the behavior of turbulent plasma flows, which are closely coupled to the magnetic fields in the corona.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

Turbulent Plasma Motion and Energy Dissipation: Turbulent motion in the corona is another mechanism that contributes to coronal heating. In the turbulent regime, energy is transferred from large-scale structures to smaller scales through **eddy dissipation**, increasing the local temperature. The turbulent energy dissipation rate can be described by: $\varepsilon = \nu * (\partial u / \partial x)^2$, where **ν** is the kinematic viscosity of the plasma, **u** is the velocity of turbulent motions and $\partial u / \partial x$ is the spatial gradient of the velocity field. The turbulent cascades in the corona can dissipate magnetic energy, leading to localized heating of the plasma. Current-driven instabilities like **kink instabilities** in twisted magnetic loops also can enhance turbulence and contribute to energy dissipation. Additionally, **shear flows** in the corona, driven by magnetic fields, could further contribute to energy dissipation through viscous friction and nonlinear interactions.

Turbulent Transport Models: Turbulence in the corona can be modeled using statistical mechanics and fluid dynamics. Models of turbulent energy transport describe how energy cascades from large-scale, low-frequency waves to smaller scales, where it is dissipated and converted into heat. During turbulent times much energy can be transferred or tranformed from large, low-frequency motions to smaller, high-frequency eddies. As the turbulence dissipates on small scales, it converts mechanical energy into heat, which then raises the temperature of the surrounding plasma. This energy cascade is described by Kolmogorov-type turbulence models, which are frequently used to model turbulent flows in various astrophysical and laboratory plasmas. **The energy dissipation rate is given by:** $\varepsilon = \nu (\partial u / \partial x)^2$, where **ν** is the kinematic viscosity, **u** the velocity field of the turbulent flow and **x** is the spatial coordinate. This **Kolmogorov turbulence model** helps describe how large-scale turbulent motions in the corona break up into smaller eddies, dissipating energy as heat. As turbulent energy is transferred to smaller scales, the energy density increases, and this leads to local heating of the plasma. It is a similar example like nanoflares, who can show that many small-scale actions can have big impact in the sum.

Summary of Turbulence in Coronal Heating: Turbulence in the solar corona contributes significantly to coronal heating by promoting the dissipation of **Alfvén waves**, **magnetoacoustic waves**, and other **magnetohydrodynamic (MHD) waves**. These waves travel along the magnetic field lines and, under turbulent conditions, can experience **mode conversion** and **dissipation**, releasing their energy as heat in the corona. The interaction of turbulent eddies with magnetic fields can also generate **current sheets**, where magnetic reconnection occurs, further releasing magnetic energy and heating the corona. The key role of turbulence is not only in enhancing wave dissipation but also in increasing the rate of magnetic reconnection. This, in turn, contributes to the overall heating of the solar corona. In this sense, turbulence enhances both the **magnetic energy dissipation** and **wave-particle interactions**, both of which are essential processes for the heating of the solar corona. Understanding these turbulent processes, especially in conjunction with other mechanisms such as **magnetic reconnection**, **wave heating**, and **Alfvén wave dissipation**, is critical for resolving the long-standing mystery of the **coronal heating paradox**. While gravitational stratification and standard heat transport mechanisms (such as thermal conduction) are important, they alone cannot account for the observed extreme temperatures of the corona. The combination of major processes, including magnetic dynamic, turbulence and wave dissipation, likely work synergistically to produce the high-energy environment seen in the Sun's corona.

26. Wave-Particle Interactions and Energy Dissipation

A crucial component of understanding coronal heating lies in the interactions between waves and particles, particularly through processes like **stochastic acceleration** and **resonant wave-particle interactions**. These processes play a significant role in transferring energy from waves to particles, thereby heating the plasma in the corona. The **Fokker-Planck equation** is a key tool used to describe the evolution of the particle distribution function in the plasma, which helps model wave-particle interactions. The general form of the equation is: $\partial f / \partial t = \partial / \partial p \, (D_{pp} \, \partial f / \partial p)$, where **f** is the distribution function, D_{pp} is the momentum diffusion coefficient and **p** is the momentum of the particles.This equation describes how the particle distribution function **f** evolves in time due to various processes, including the interaction with waves.

To model the effect of solar radiation on the particles, we can modify the equation to include the impact of solar emissions: $\partial f / \partial t = \partial / \partial p \, (D_{pp} \, \partial f / \partial p) + \gamma \cdot E_{solar}$, where **γ** is a factor representing the impact of solar radiation and E_{solar} is the energy from solar emissions that interacts with the particles. These wave-particle interactions lead to a more efficient transfer of energy to the particles, contributing to their acceleration. **Resonant interactions** are particularly efficient in this regard, as they lead to the direct

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

transfer of wave energy to particles with resonant velocities, whereas **non-resonant interactions** are less efficient but still contribute by spreading energy over a broader range of particle velocities.

Wave-particle interactions in the solar corona can also be modeled using **kinetic theory**, where the distribution of particles in phase space is governed by the **Boltzmann transport equation**. In the context of a plasma, this equation describes how the distribution function **f(v, r, t)** evolves over time, considering particle motion and external forces: $\partial f / \partial t + v \cdot \nabla f + (F/m) \cdot \nabla_v f = C[f]$, where **f** is the distribution function of particles (in phase space), **v** the particle velocity, ∇ the gradient operator in real space, ∇_v the gradient operator in velocity space, **F** the force acting on the particles (e.g., electromagnetic force), **m** the mass of the particle and **C[f]** is a collision term representing the interactions of particles with each other or with wavesThe collision term **C[f]** accounts for the interactions such as **wave-particle interactions**, **scattering**, and **particle collisions**, which dissipate energy and convert the energy carried by waves into heat. These interactions are fundamental to the heating process, as they provide the mechanism by which the energy in waves is transferred to the particles in the corona, raising their temperature.

High-Frequency Waves in Coronal Heating: One of the key unresolved issues in solar physics is understanding the role of **high-frequency waves** in the coronal heating process, especially in relation to **Alfvén waves** and **fast-mode magnetoacoustic waves**. While **low-frequency Alfvén waves** have been well studied, the behavior of higher-frequency waves is still uncertain, though they are thought to be crucial for efficiently dissipating energy in the corona. These high-frequency waves are difficult to detect and model due to their complex interaction with the solar plasma. Recent advances in solar observational instruments, including those on the **Parker Solar Probe (PSP)**, have provided new insights into high-frequency waves and their potential role in energy transfer. Observations indicate that **Alfvén waves** are more prevalent in the **solar wind** than previously thought, with the **FIELDS instrument** aboard the Parker Solar Probe detecting high-frequency oscillations that may play a significant role in wave dissipation and energy transfer to the corona. Other instruments, such as the **Helioseismic and Magnetic Imager (HMI)**, **SDO (Solar Dynamics Observatory)**, and **Hinode**, have captured important data on wave-particle interactions. For example, **Alfvén waves** propagating from the **photosphere** into the corona have been observed in **coronal loops** and **active regions**, with these oscillations correlating with localized heating. These findings reinforce the idea that wave dissipation is a critical component of coronal heating.

Wave-Particle Interactions and Heating Observations: Several solar missions have provided valuable data that help connect wave-particle interactions to coronal heating. Observations from instruments aboard **Hinode**, **IRIS (Interface Region Imaging Spectrometer)**, and **SDO** have confirmed that wave-particle interactions in the corona contribute to localized heating. **Hinode's EUV Imaging Spectrometer (EIS)**, for example, has observed high-energy tails in the electron population, indicative of **non-thermal distributions** produced by wave heating. Similarly, **IRIS** has provided data showing turbulent motions in the transition region, which are known to enhance wave-particle interactions and contribute to coronal heating. In addition, the **Advanced Composition Explorer (ACE)** spacecraft has detected high-energy particles, such as protons and electrons, which are accelerated during solar flare events. Many of these particles are likely the result of wave-particle interactions within the corona, further supporting the link between wave dissipation and coronal heating.

The Role of High-Energy Particles in Coronal Heating: The detection of **high-energy particles** in the solar wind and during **solar flare events** also highlights the importance of wave-particle interactions. These energetic particles, particularly protons and electrons, are often accelerated in the corona through resonant wave-particle interactions, contributing to the observed heating of the plasma. The **solar wind** is thought to be influenced by such interactions, which are believed to help sustain the high temperatures of the corona. Furthermore, during **coronal mass ejections (CMEs)** and **solar flares**, high-energy particles are generated, many of which are the result of wave-particle interactions. These high-energy particles further contribute to the heating and dynamics of the corona. The acceleration of these particles, along with their interaction with waves in the corona, plays a crucial role in the overall energy balance and temperature distribution of the solar atmosphere.

27. Wave Types, Propagation and Energy Transport

Alfvén waves, **magnetoacoustic waves**, and **slow magnetoacoustic waves** play a key role in transporting energy from the lower layers of the Sun's atmosphere (like the photosphere) to the corona.

Alfvén Waves are oscillations in the magnetic field of the plasma and are thought to carry energy from the solar interior to the corona. They propagate along magnetic field lines. The **dispersion relation** for Alfvén waves in the corona is: $\omega^2 = k^2 v_A^2$, where **ω** is the wave frequency, **k** the wave vector and **v_A** is the Alfvén speed.

A hybrid wave that bridges the gap between Alfvén waves and ion-acoustic waves **are Kinetic Alfvén Waves (KAWs)**. They involve wave motion that is highly influenced by the kinetic properties of ions and electrons in the plasma. KAWs are a form of Alfvén waves in higher-order that involve **kinetic effects** rather than just **magnetohydrodynamic (MHD)** effects. They occur at scales smaller than the ion inertial length. These waves propagate with **perpendicular electric fields** to the magnetic field lines and have a characteristic wavelength that depends on the plasma parameters. The interaction of KAWs with electrons and ions can cause **stochastic heating** and energy dissipation at smaller scales. Kinetic Alfvén Waves are proposed to contribute to **coronal heating** by transferring wave energy into electrons and ions through nonlinear interactions, such as wave-particle resonance. These waves are believed to play a role in **small-scale heating** in the corona and could help explain the **high temperatures** of the **coronal loops**.

Another type of **magneto-acoustic waves** are **Kink Waves**, also known as Coronal Loops Oscillations. They can occur in coronal loops, where the magnetic field supports the oscillatory motion of plasma along the loop. These waves are named after the "kink" that can be observed when a coronal loop oscillates or vibrates, causing the loop's shape to distort in a characteristic manner. They are important for understanding the **dynamics** of **solar coronal loops** and their role in transferring energy through the corona. Kink waves involve the motion of plasma along the magnetic field lines of a coronal loop. When excited, these waves cause a displacement in the plasma, resulting in a periodic **bending** or **kinking** of the loop. The wave can propagate as a global oscillation, with the plasma oscillating back and forth, driven by forces such as magnetic tension and plasma inertia. The frequency of a kink wave depends on the magnetic field strength, the loop length, and the plasma density. The resonant frequency of the kink wave is typically lower than that of other magneto-acoustic waves. The frequency of kink oscillations in a coronal loop can be approximated by: $\omega = (2\pi) * (v_A / L)$, where **v_A** is the Alfvén speed and **L** the loop length. Kink waves can play a significant role in **energy transfer** within the solar corona. These waves can carry **kinetic energy** and, when they reach higher altitudes in the corona, they may dissipate through processes such as **wave damping** or **resonance** with other wave modes, such as **acoustic waves** or **Alfvén waves**. As kink waves propagate along coronal loops, their energy may get dissipated due to **non-linear interactions**, causing local heating of the plasma. The magnetic tension that confines the wave may also cause **magnetic reconnection** or **wave coupling**, which can lead to **localized heating** and the production of **high-energy particles**. In certain circumstances, kink waves may trigger **magnetic reconnection events**, which are thought to be a major source of coronal heating. These events release vast amounts of energy, which could help explain the high temperatures observed in the corona. The energy transfer from kink waves to the plasma can be described by the **dissipation rate** due to **wave-particle interactions**: $P = (\omega^2 B^2 / 8\pi) * (1 / k^2)$ where **ω** is the wave frequency, **B** the magnetic field strength and **k** is the wave number. This formula indicates how the **magnetic energy** in the wave can be transferred into the plasma as heat.

Electron-Cyclotron Waves (ECWs) are a type of wave that arises in electromagnetic fields and magnetized plasmas. The can occur especially in the presence of strong magnetic fields, such as those found in the solar atmosphere, planetary magnetospheres, and laboratory plasma systems. The waves are closely tied to the behavior of electrons within the magnetic field, where they undergo **cyclotron motion**. If electrons in a plasma move in circular or helical paths along magnetic field lines, this movement can create oscillations at the **electron cyclotron frequency**, denoted as: $\omega_e = eB / m_e$, Where **e** is the electron charge, **B** the magnetic field strength and **m_e** is the electron mass. The interaction between the electron motion and the electromagnetic fields at this frequency generates **ECWs**. These waves are electromagnetic in nature and can couple with other types of plasma waves, such as ion-cyclotron waves or magnetoacoustic waves, to transfer energy and cause heating, particularly in regions with high magnetic field strengths. The frequency of electron-cyclotron waves is dependent on the magnetic field strength and can range from kilohertz to gigahertz, making them a type of **high-frequency wave** in the solar atmosphere. ECWs can propagate along magnetic field lines, and their behavior is highly dependent on both the electron density and the magnetic field configuration in the plasma. These waves typically interact with the plasma in a way that can lead to wave-particle interactions, which are critical in understanding energy transfer processes in the corona. Electron-cyclotron waves have been identified as playing a key role in **coronal heating** due to their ability to interact with high-energy electrons. As these waves propagate through the solar corona,

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

they can dissipate their energy via wave-particle interactions. This dissipation occurs when the energy from the ECWs is transferred to resonant electrons, accelerating them. These high-energy electrons can then collide with other particles in the corona, transferring their energy and leading to localized heating. The interaction of ECWs with electrons, particularly through **Landau damping** and other non-linear interactions, plays a significant role in how the wave energy is absorbed by the plasma. Landau damping occurs when waves interact with resonant particles (electrons) whose velocities match the phase velocity of the wave, transferring energy from the wave to the particles, causing them to accelerate. The accelerated electrons then contribute to the heating of the plasma by colliding with ions, increasing the overall temperature of the corona.

Fast and Slow Magnetoacoustic Waves: The fast waves propagate along **magnetic field lines** and can efficiently heat the plasma, especially in regions of high magnetic activity. They interact with both **electrons** and **ions** in the corona, causing them to accelerate and gain energy through collisions and resonance with the wave. **Fast-mode waves** are responsible for the **global heating** of the corona, as they can propagate over long distances and deposit energy into both low and high-atomic-number elements such as **H**, **He**, **O**, and **Fe**. **Slow magnetoacoustic waves** have a slower propagation speed and are often associated with localized heating near coronal loops and sunspots. While slower than fast-mode waves, slow-mode waves still contribute to **local heating** by exciting **resonant ionospheric modes**, which leads to energy transfer to specific ions like $\mathbf{O^{6+}}$ and $\mathbf{Ne^{6+}}$. These waves are particularly important in determining the fine structure and temperature variation in the **coronal loops**. The interaction of the solar corona with both **Alfvén waves** and **magnetoacoustic waves** plays a central role in the **energy dissipation** and **heating processes** that shape the **dynamic environment** of the Sun's outer atmosphere. Observations from the **HMI** on the **SDO** reveal how **magnetoacoustic waves** travel upward from the Sun's surface and are likely responsible for **coronal heating** by depositing energy in the form of **shocks** and **compression** as they propagate through the corona. The observations suggest that **wave-driven heating** occurs more efficiently than initially thought, with acoustic waves acting as a precursor for **magnetic energy transfer** to higher altitudes, further fueling the thermal energy in the corona. Fast magnetoacoustic waves tend to be more effective at higher frequencies, whereas slow waves are important in the lower parts of the corona. These waves can be dissipated in the corona, leading to the deposition of energy and heating of the plasma.

The dispersion relation for a magnetoacoustic wave in a magnetized plasma is: $\omega^2 = k^2 c_s^2 + k^2 v_A^2$, where ω is frequency of the wave, k the wave number, c_s the sound speed and v_A is the Alfvén speed.

The sound speed is given by: $c_s^2 = (\gamma p / \rho) + (B^2 / (\mu_0 \rho))$, where γ is adiabatic index and the ratio of specific heats, p the plasma pressure, ρ the plasma density, B the magnetic field strength and μ_0 is the permeability of free space. In this equation, c_s represents the sound speed in the plasma, which depends on both the pressure p and the magnetic field B. Magnetoacoustic waves combine the effects of thermal pressure and magnetic fields, and their dissipation can lead to energy deposition in the corona, contributing to the observed high temperatures.

Magnetosonic waves, which combine both compressive and Alfvén-like characteristics, also contribute to heating by **compressing plasma** as they propagate through the corona. These waves may also interact with density fluctuations and undergo **nonlinear steepening**, which significantly amplifies the wave's energy. The energy flux associated with magnetosonic waves can be expressed: $F_{ms} = (\rho * V * E_{wave}) / (L_{wave})$, where ρ is the plasma density, V the wave velocity, E_{wave} the wave energy and L_{wave} is the characteristic length of the wave. Empirical data from **coronal seismology** indicate the presence of compressive waves in coronal loops, further supporting the role of these waves in heating mechanisms. Numerical simulations have shown that nonlinear interactions between different MHD waves can lead to efficient energy dissipation and provide a viable solution to the coronal heating paradox.

Fast magnetosonic waves and **slow magnetosonic waves** are fundamental to energy transport in the solar corona. These waves involve both plasma compression and magnetic field perturbations. The fast waves are **compressional waves** that travel with a speed greater than the **Alfvén speed** and involve both the plasma and magnetic field. They can dissipate energy through **magnetic compression** and **thermal conduction**, contributing to heating in the corona. They can also couple with **fast-mode shocks** in the corona, leading to energy dissipation in the form of **shock heating**. Slower waves as Alfvén waves can be more easily damped in the lower corona due to their interaction with plasma density and temperature gradients. Slow magnetosonic waves are considered to be important for **energy dissipation** and **wave heating** through processes like **Landau damping** and **resonant absorption**. The dispersion relation

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

for these waves in a magnetized plasma is given by: $\omega^2 = k^2 v_A^2 + k^2 c_s^2$, where **v_A** is the **Alfvén speed** and **c_s** is the **sound speed**.

The fast and slow magnetosonic waves mentioned above are often referred to as **fast mode** and **slow mode** in the broader context of MHD theory. These waves travel through the solar atmosphere and play crucial roles in energy dissipation and **coronal heating**. The fast mode is a compressive wave that propagates faster than both the Alfvén and slow modes. It can result in **shock formation** and **heating**, particularly if interacting with density gradients in the corona. The energy density in a **fast magnetosonic wave** is given by the sum of the kinetic and magnetic components: $W_{fast} = \frac{1}{2}\rho v^2 + B^2 / (2\mu_0)$, where **ρ** is the plasma density, **v** is the velocity of the wave, and **B** is the magnetic field strength. The **slow mode** is the more **subsonic** counterpart, typically propagating in regions where the plasma is less dense. It is more sensitive to **thermal gradients** and **density variations** in the corona, leading to energy deposition and local heating.

Ion-cyclotron waves can play an important part in coronal heating processes. It is a type of low-frequency electromagnetic wave that propagate in magnetized plasmas. These waves are primarily associated with the **gyro-motion of ions around magnetic field lines** and are important for energy transfer and heating in the solar corona. They are particularly relevant in regions where the ion temperature is high, as they can resonate with the motion of ions, **leading to efficient energy dissipation and plasma heating**. Ion-cyclotron waves are generated when ions in a plasma resonate with the electromagnetic wave's frequency, leading to energy absorption. These waves propagate perpendicular or obliquely to the magnetic field lines and can transfer energy through **ion-ion collisions** and **wave-particle interactions**. The resonant frequency of ion-cyclotron waves depends on the charge-to-mass ratio of the ions and the strength of the magnetic field: $\omega_c = (q * B) / m$, where **q** is the ion charge, **B** the magnetic field strength and **m** the ion mass. In the solar corona, the presence of multiple ion species can lead to a spectrum of cyclotron frequencies, **contributing to a broad range of energy dissipation processes**. When the wave frequency matches the cyclotron frequency of the ions, the wave energy is resonantly absorbed by the ions, leading to their acceleration and heating. This mechanism is highly efficient in low-density plasmas, such as the corona. Ion-cyclotron waves preferentially heat ions perpendicular to the magnetic field lines, leading to anisotropic temperature distributions. This effect is often observed in the solar corona, where ions exhibit higher temperatures perpendicular to the magnetic field compared to parallel directions.

The power dissipated by ion-cyclotron waves through resonant interactions can be approximated by: $P = (q^2 * B^2 / 2 * m) * v_{perp}^2$, where **v_perp** is the perpendicular velocity of ions due to wave interactions and **B** the magnetic field strength. Ion-cyclotron waves dissipate their energy through mechanisms like Landau damping (wave-particle resonance) or cyclotron resonance, where ions absorb the wave energy, resulting in local plasma heating. In turbulent plasma environments, ion-cyclotron waves can cascade to smaller scales, transferring energy to higher-frequency modes and enhancing dissipation. Ion-cyclotron waves are thus a key component of wave-based coronal heating models, helping to explain the extreme temperatures observed in the solar corona relative to the solar surface. ^^
Parker Solar Probe and **Solar Orbiter** have both made measurements indicating the presence of high-frequency waves, specifically **ion-cyclotron waves**, in the outer solar atmosphere. These waves are thought to transfer energy to the **ionized particles** in the corona by resonating with their gyromotion. Data from **solar wind measurements** near the corona indicate that ion-cyclotron resonance could be responsible for accelerating ions in the solar wind, which in turn heats the corona. The **resonant absorption of wave energy** in the corona by ions is thought to be a significant energy source, contributing to the **high temperatures observed** in the solar corona. This process is thought to be particularly effective in regions of the solar atmosphere with strong magnetic fields.

Langmuir waves are high-frequency **electron plasma oscillations** in a plasma, where the oscillations involve the collective motion of electrons around a mean position. These waves play a significant role in many astrophysical environments, including the solar corona, where they contribute to the **energy transfer** and **particle acceleration** in regions of intense magnetic activity. Langmuir waves are generated when the electron density fluctuations in the plasma become coherent and oscillate with a frequency much higher than the ion plasma frequency. These waves are associated with the high-frequency oscillations of the electron plasma and are often coupled with lower-frequency ion waves. **The frequency of Langmuir waves is given by:** $\omega_L = \sqrt{(4\pi e^2 n_e / m_e)}$, where **e** is the electron charge, n_e the electron density and m_e the electron mass. This is a characteristic frequency that depends on the local electron density, and it is typically much higher than the **cyclotron frequency** of electrons. Langmuir waves can interact with other plasma waves, such as **electron-cyclotron waves** and **acoustic waves**, and contribute to **energy**

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

dissipation. The wave-particle interactions can accelerate electrons, which can then collide with surrounding ions, transferring their energy and resulting in **heating of the coronal plasma**. Langmuir waves may experience **non-linear decay**, where their energy is transferred to **lower-frequency waves** or is dissipated through **electron Landau damping**, where the wave energy is absorbed by resonant electrons. The power dissipated by **Langmuir waves can be modeled by: $P = (n_e e^2 / m_e) * (\delta v)^2$** , where **$\delta v$** is the velocity fluctuation of the electrons due to wave-particle interactions.

Whistler Waves and Plasma Dynamics: These waves can propagate through plasma in the corona and to outer regions of the magnetosphere. They are often associated with high-frequency electromagnetic disturbances and are typically found at high altitudes in the solar atmosphere. The waves are significant in explaining some of the fine structures and temperature variations observed in the solar corona, as well as playing a key role in electron heating, which is an essential factor in the ongoing investigation of the coronal heating paradox. They are typically in the **lower hybrid resonance frequency range** or **electron cyclotron frequency range**, depending on their specific environment. The propagation of whistler waves is closely related to the dynamics of the plasma, particularly the interactions between electrons and the magnetic field. In a plasma environment like the Sun's corona, the electrons are more responsive to these waves due to their lower mass compared to ions, leading to the electron heating. Whistler waves are especially important because they can interact with electrons in such a way that they provide an additional mechanism for **coronal heating** beyond traditional mechanisms like **Alfvén waves** and **magnetoacoustic waves**.

Whistler waves are defined by their **polarization**, which is typically **left-hand circular polarization** in the plasma. These waves propagate along the **magnetic field lines** and are often observed in regions with **high electron densities** and strong magnetic fields. The dispersion relation for whistler waves is given by: **$\omega^2 = \omega_pe^2 + \omega_ce^2 / (1 + \omega_ce / \omega)$** , where **$\omega$** is the wave frequency, **ω_pe** is the plasma frequency with $\omega_pe = \sqrt{(n_e * e^2 / (\varepsilon_0 * m_e))}$ and the electron cyclotron frequency is **$\omega_ce = eB / m_e$**. The term **n_e** stands for the electron number density, **e** is the elementary charge, **B** the magnetic field strength, **ε_0** the permittivity of free space and **m_e** is the mass of the electron. This relationship demonstrates how the whistler wave frequency depends on both the plasma density and the magnetic field strength. As these waves propagate, they can undergo various **wave-particle interactions**, and their energy can be transferred to the plasma, resulting in **electron acceleration** and **heating** - which is also one of the primary reasons they are relevant to the **coronal heating paradox**. The energy transferred from the wave to the **electrons** can cause the electrons to move to **higher energy states**, thus increasing the overall **thermal energy** of the plasma. This electron heating can significantly contribute to the high temperatures observed in the solar corona. Wave-particle interactions that lead to **electron acceleration** include **resonant interactions** between the electron's gyrofrequency and the wave frequency. When the wave frequency matches the **cyclotron frequency** of the electron, the electron can absorb energy from the wave, leading to an increase in its kinetic energy. **Whistler waves** interact with **electron populations** in the form of **stochastic acceleration**, where the interaction between electrons and the fluctuating electromagnetic fields of the waves results in a **non-thermal energy distribution**. This is crucial because it not only heats the electrons directly but can also contribute to the generation of high-energy tails in the electron distribution, which have been observed in the solar corona and are thought to play a role in the observed X-ray emissions.

The **spatial distribution** and **frequency spectrum** of whistler waves in the corona could provide insight into the **fine-scale temperature structures** of the solar atmosphere. The existence of high-frequency whistler waves could be associated with localized heat deposition in small regions of the corona, which might explain some of the **temperature inhomogeneities** and **temperature spikes** observed in **coronal loops**. The dissipation of whistler wave energy leads to enhanced electron temperatures in regions where the waves are most intense. As the whistler waves travel upward through the **coronal magnetic field**, they could contribute to the **heating of plasma structures** along magnetic field lines, possibly explaining the **coronal heating paradox** by directly addressing the issue of how energy is transferred to the corona from lower layers. Whistler waves can also play an important role in the **acceleration of the solar wind**. These waves are often observed in the **outer corona** and are thought to contribute to the dynamics of the **solar wind**. They could influence the **distribution of particles** in the corona and provide additional energy that drives the **outflow** of charged particles from the Sun, helping to explain the high-speed solar wind observed in certain regions such as **coronal holes**. More about CMEs, MHD, particle and plasma interactions with different wave types will be released in the second edition of the Sun's Water Theory - including advanced research about further problems and solutions.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

28. Wien's Law and the Coronal Heating Paradox

Wien's Displacement Law provides a relationship between the temperature of an object and the wavelength at which it emits radiation most efficiently. It is a crucial piece in understanding the spectral properties of stars, including the Sun, and helps explain some of the key features of the coronal heating paradox. To delve deeper into this, we need to explore both the theoretical and practical implications of this law, particularly how it links the temperature of the photosphere to the observed spectrum, and how it can be used in the context of the Sun's corona. **Wien's Displacement Law can be expressed mathematically as:** $\lambda_{max} = b / T$, where λ_{max} is the **peak wavelength** (in meters), **b** is the **Wien's constant** with 2.898×10^{-3} m·K, and **T** is the temperature of the blackbody in Kelvin (K). This law tells us that the peak wavelength of emission is inversely proportional to the temperature of the object. For the Sun, with a temperature around 5,800 K at the photosphere, the peak wavelength of emitted radiation is around **500 nm**, which corresponds to **green-yellow light** in the visible spectrum.

Applications in Coronal Heating Models: In recent research, understanding **Wien's Displacement Law** has played an important role in the development of more accurate models of coronal heating, especially in **wave heating models.** By incorporating wave energy deposition and dissipation, these models predict a temperature gradient from the photosphere to the corona that aligns with observed wavelengths. In **magnetic reconnection models**, reconnection events lead to the release of high-energy particles and radiation, which corresponds to the X-ray emission observed in the corona. These processes elevate the temperature of the corona significantly above the photosphere, causing the peak wavelength of the corona to shift well into the ultraviolet and X-ray regions, as predicted by Wien's Law. **Spectral energy distribution models** of the Sun's atmosphere, including the photosphere, chromosphere, and corona, use Wien's Law as a tool to refine their predictions about the radiative output from these layers. By considering the temperature at each layer and applying Wien's Law, researchers can estimate the **spectral energy distribution (SED)** and refine models that simulate the Sun's observed emissions across the entire electromagnetic spectrum.

Bridging the Paradox: Wien's Displacement Law is a powerful tool in understanding the spectral properties and frequencies of the Sun. It provides essential insights into the coronal heating paradox. By applying this law to the Sun's atmosphere, it is clear that the corona emits primarily in the UV and X-ray regions, despite being much hotter than the photosphere. This **shift in the peak wavelength** is due to the **non-equilibrium processes** that govern the energy transport in the corona, such as **wave heating**, **magnetic reconnection**, and **turbulent dissipation**. The temperature discrepancy between the photosphere and the corona is resolved not by classical heat conduction, but by non-radiative energy input from the lower atmosphere, magnetic field dynamics, and wave-driven heating mechanisms. The law serves as a foundational principle for understanding why the corona's emission peaks in the short wavelength regions and thus provides an essential piece of the puzzle in explaining the high temperatures observed in the Sun's outer atmosphere.

The application of Wien's Law also helps refine models of **energy transport** in the corona, as it provides a direct relationship between temperature and the emitted radiation. **SOHO's CDS** and **STEREO's COR1** have contributed spectroscopic data that align with this model, revealing that the coronal spectrum is dominated by high-energy wavelengths that are consistent with the high temperatures measured in the corona. Furthermore, **Hinode's EIS** has provided important spectroscopic data that show **Doppler shifts** and **line broadening**, consistent with the heating processes that occur due to wave-particle interactions and magnetic reconnection. These features support the idea that magnetic activity and wave propagation are responsible for the high temperatures observed in the corona, as predicted by the shift in wavelength. **Hinode's XRT** and **SDO's AIA** have observed this transition in the solar spectrum, with emissions in the UV and X-ray regions becoming increasingly prominent as the temperature rises in the corona. The X-ray emissions detected by these instruments directly correlate with the high temperatures observed in the corona, suggesting that the Sun's outer atmosphere is heated by non-radiative processes.

The Wien's Displacement Law is essential for understanding why the **photosphere** emits primarily in the visible spectrum and with the peak wavelength, a direct consequence of the surface temperature (~5,800 K). The relationship can help to **quantify the distribution of energy** across the electromagnetic spectrum based on temperature. The law also highlights an important point in the context of **coronal heating**. It shows that for a higher temperature (such as that in the corona, which can exceed **1-2 million K**), the peak wavelength of emission would shift to **much shorter wavelengths** (in the **X-ray** and **ultraviolet** ranges). This is why we observe the **corona's emission** primarily in the **ultraviolet (UV)** and **X-ray regions**.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

even though these wavelengths are far from the Sun's peak emission in the visible. This shift is significant for understanding how the **corona** can reach temperatures far exceeding those of the **photosphere**.

Research Data from Upcoming Solar Missions

With upcoming missions like **Chromospheric Lyman-Alpha Spectro-Polarimeter (CLASP2)** and the **Extreme Ultraviolet Spectroscopic Telescope (EUVST)** set to launch in the next few years, a wealth of new observational data is expected to refine our understanding of coronal heating. These missions will provide high-resolution spectroscopic and imaging data across the solar atmosphere, including both the chromosphere and the corona. Key research areas are **high-resolution observations** of the **chromospheric magnetic fields** and their influence on **coronal heating through wave propagation**. Improved measurements of particle acceleration and heating during solar flares could provide direct evidence of how magnetic reconnection events contribute to heating on both global and local scales. Observation of coronal loops in different wavelengths (e.g., UV and X-rays) help to study the **dynamics of energy transport** and how energy is transferred to the plasma from the lower layers of the atmosphere. Data from these missions could lead to **new insights** into the roles of **magnetic fields**, **waves**, and **plasma interactions** in the **coronal heating process**. It may also provide better observational constraints for modeling the solar corona and help resolve existing theoretical uncertainties. The new **observational findings** and **empirical evidence** presented in this article significantly advance our understanding of the **coronal heating paradox**. The contributions from all the processes and mechanisms, which are explained in numerous sections, point to a **multi-faceted** approach to understanding the heating of the Sun's outer atmosphere. Some aspects still remain unclear, such as the precise **role of nanoflares**, the exact **distribution of heating** in the corona, and how **magnetic fields** and **plasma flows** interact at various scales. **Future missions**, such as **CLASP2** and **EUVST**, will be crucial for providing more **detailed and high-resolution observations** of the solar atmosphere, and will offer the empirical data needed to refine existing models and test new hypotheses.

References and Inspirational Sources:

- **Allard, F., et al. (1997).** "Structure and Temperature of the Solar Photosphere." *Astrophysical Journal*, 484(2), 623-640.
- **Bradshaw, S. J., & Cargill, P. J. (2013).** A Review of Solar Coronal Heating. *Space Science Reviews*, 175(1-4), 3-29.
- **Burlaga, L. F. (1991).** "Bow Shocks and Their Relevance to Solar Wind Interactions." *Journal of Geophysical Research*, 96(A1), 1-13.
- **Chen, P. F., et al. (2006).** "Radiative Cooling in Solar Flares and the Heating of the Coronal Plasma." *Astrophysical Journal*.
- **Chitta, L. P., et al. (2021).** "Magnetic Reconnection and Wave Heating in the Solar Corona: New Insights from Parker Solar Probe." *Astrophysical Journal*, 921(1), 59.
- **Chou, D.-Y., & Chou, C.-H. (2001).** "Opacity and Energy Transport in the Solar Atmosphere." *Solar Physics*, 199(2), 167-182.
- **Cohen, L. M., et al. (2022).** "Magnetic turbulence in the solar corona:" New insights from Parker Solar Probe and Solar Orbiter. *Astrophysical Journal*, 937(2), 145-159.
- **Cranmer, S. R. (2009)**. "Coronal Heating: The Role of Turbulence." *Solar Physics*, 255, 125-145.
- **De Pontieu, B., et al. (2020).** Wave-driven heating in the solar corona. *Solar Physics*, 295(4), 50-67.
- **De Pontieu, B., et al. (2023).** "Coupling Between the Chromosphere and Corona: New Insights from ALMA and IRIS." *Astrophysical Journal*, 911(2), 215.
- **Forbes, T. G. (2000).** "The Solar Wind and Bow Shocks." *Space Science Reviews*, 91(1), 17-25.
- **Gibson, S. E. (2011).** "The Solar Coronal Boundaries." *Space Science Reviews*, 158(1), 23-52.
- **Goossens, M., & Van Doorsselaere, T. (2013).** "Wave Heating in the Solar Corona." *Space Science Reviews*, 175(1-4), 59-82.

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

- **Hasegawa, A. (1994).** "Plasma Kinetic Theory and Wave-Particle Interactions." *Physics of Fluids*, 7(5), 987-995.
- **Klein, K.-L., et al. (2024).** "Role of ion-cyclotron resonance in coronal heating." *Astrophysical Journal*, 938(1), 120-133.
- **Klimchuk, J. A. (2006).** "On the Nature of Coronal Heating." *Solar Physics*, 234(1), 41-58.
- **Kuhn, J. R., & Reitzel, D. (2000).** "The Temperature Structure of the Sun's Photosphere." *Astrophysical Journal*, 541(1), 65–75.
- **Krucker, S., et al. (2022).** "The role of nanoflares in the solar corona: New findings from Parker Solar Probe." *Nature Communications*, 13(1), 2561.
- **Landi, E., et al. (2020).** "Energy Dissipation through Turbulence in the Solar Corona." *Astronomy & Astrophysics*, 636, A109.
- **Linsky, J. L. (1998).** "The Solar Atmosphere and Its Energy Balance." *Annual Review of Astronomy and Astrophysics*, 36(1), 509-535.
- **Martínez-Sykora, J., et al. (2021).** "Magnetic Reconnection and Energy Transfer in the Corona: High-Resolution Observations from DKIST." *Nature Astronomy*, 5(3), 296-304.
- **Matthaeus, W. H., et al. (2023).** "Alfvén waves and their role in coronal heating: Insights from Parker Solar Probe." *Nature Physics*, 19(6), 635-649.
- **McIntosh, S. W., et al. (2021).** "Turbulence and magnetic reconnection in the corona." *Nature Astronomy*, 5(2), 127-132.
- **NASA Solar Orbiter Team.** "Magnetic Field Interactions and Wave Propagation in the Solar Corona." *Nature Astronomy*, vol. 5, no. 3, 2021.
- **Okamoto, T. J., et al. (2016).** "Observing the Sun with the Solar Dynamics Observatory." *Space Science Reviews*, 204(1-4), 95-133.
- **Parker, E. N. (1988).** "Nanoflares and the Heating of the Solar Corona." *Astrophysical Journal*, 330, 474-479.
- **Pesnell, W. D., et al. (2022).** "The Role of the Solar Transition Region in Coronal Heating." *Solar Physics*, 297(9), 120.
- **Priest, E. (2014).** "The Physics of the Sun: A University Introduction." *Cambridge University Press.*
- **Kulsrud, R. M. (2005).** "Plasma Physics for Astrophysics." *Princeton University Press.*
- **Roberts, B. (2000).** "Wave Propagation in the Solar Atmosphere." *Solar Physics*, 193(1), 9-28.
- **Saha, K. (2011).** "The Total Energy Output of the Sun and Its Variations." *The Astrophysical Journal.*
- **Spruit, H. C. (1999).** "Gravity and the Solar Coronal Stratification." *Astronomy & Astrophysics*, 349(3), 982-994.
- **Spruit, H. C. (2006).** "The Dynamics of Solar Convection and Coronal Heating." *Space Science Reviews*, 125(1-3), 171–194.
- **Tian, H., et al. (2023).** "Coronal heating through wave propagation and energy dissipation: Spectroscopic evidence from SDO. *Solar Physics, 296(4), 45-63.*"
- **Thomas, R. J., & Montesinos, B. (1991).** "Wien's Law and Coronal Heating." *Nature*, 353, 825-828.
- **Toriumi, S., & Yokoyama, T. (2017).** "Magnetic Reconnection and Solar Flares." *Solar Physics.*
- **Turner, L. D., & Cally, P. S. (2017).** "Plasmonic Heating in Solar Coronal Plasma." *Astrophysical Journal*, 843(2), 175.
- **Vernazza, J. E., Avrett, E. H., & Loeser, R. (1981).** "Structure of the Solar Chromosphere. III—The Semiempirical Model of the Solar Chromosphere." *Astrophysical Journal*, 45, 635-642.
- **Verwichte, E., et al. (2012).** "MHD Waves and Their Role in Heating the Solar Corona." *Living Reviews in Solar Physics.*

 This artistic and scientific work is protected under national and international laws. Unauthorized reproduction, scanning, distribution, digital processing and copying is strictly prohibited without written consent from the author. Read only and fair use.

- **Wang, T., et al. (2022).** "Wave-Driven Heating and Solar Wind Acceleration: A New Perspective from Parker Solar Probe." *Astrophysical Journal Letters*, 924(1), L15.
- **Wu, S. T., & Lee, L. C. (1979)**. "The Role of Magnetic Reconnection in Solar Wind Acceleration."
- **Yamada, M., et al. (2010)**. "Magnetic Reconnection and Wave-Particle Interactions." *Nature Physics*, 6(4), 176-180.
- ***Zhao, L., et al. (2023)***. "Coronal bright points: A closer look at localized heating mechanisms." *Solar Physics, 298(5), 211-222.*

More references, sources and links will come into the final publication and after adequate funding. In the next publication will follow a short executive summary of some major key factors and selected topics. The article is mainly for research and further improvements of the study.

This pre-publication is an extract of the scientific paper for the advanced edition of the Sun's Water Theory and study. The research project is focused on solar physics, solar science and sunlight research. This article includes also parts of the new book called **"The Corona Paradox"**, which will be released in spring 2025. It will be a comprehensive artwork about the scientific problems and potential solutions, including practical and theorethical solution approaches, as well as great examples and many references – similar like here and in the previous studies and scientific writings. Copies are not allowed without the permission of the author. The document was created on basis of the manuscript and on one final draft version.

This document, including its **artistic, private, operational, scientific, and sensitive content**, is protected under **national** and **international laws,** including but not limited to the **Berne Convention (1886)**, **Copyright Laws**, the **Digital Millennium Copyright Act (DMCA)**, and the **WIPO Copyright Treaty (1996). All rights are reserved by O.G.Caplikas**, the author of **Suns Water Theory and study**. **No part of this document - whether text, artwork, or other proprietary content - may be copied, digitized, photographed, processed, scanned and / or shared in any similar form** without explicit written authorization from the author. Unauthorized use can result in legal consequences. The document includes important research information, discoveries and data which is protected by several protection rights. There are unique creations and certain combinations were not published before. Some texts will be used in further publications, which will be released only as print version – this is also one reason for this disclaimer.

This document, including its **artistic, private, operational, scientific, and sensitive content**, is protected under **national** and **international laws,** including but not limited to the **Berne Convention (1886)**, **Copyright Laws**, the **Digital Millennium Copyright Act (DMCA)**, and the **WIPO Copyright Treaty (1996). All rights are reserved by O.G.Caplikas**, the author of **Suns Water Theory and study**. **No part of this document - whether text, artwork, or other proprietary content - may be copied, digitized, photographed, processed, scanned and / or shared in any similar form** without explicit written authorization from the author. Unauthorized use can result in legal consequences. The document includes important research information, discoveries and data which is protected by several protection rights. There are unique creations and certain combinations were not published before. Some texts will be used in further publications, which will be released only as print version – this is also one reason for this disclaimer.

This document, including its **artistic, private, operational, scientific, and sensitive content**, is protected under **national** and **international laws,** including but not limited to the **Berne Convention (1886)**, **Copyright Laws**, the **Digital Millennium Copyright Act (DMCA)**, and the **WIPO Copyright Treaty (1996). All rights are reserved by O.G.Caplikas**, the author of **Suns Water Theory and study**. **No part of this document - whether text, artwork, or other proprietary content - may be copied, digitized, photographed, processed, scanned and / or shared in any similar form** without explicit written authorization from the author. Unauthorized use can result in legal consequences. The document includes important research information, discoveries and data which is protected by several protection rights. There are unique creations and certain combinations were not published before. Some texts will be used in further publications, which will be released only as print version – this is also one reason for this disclaimer.